

FLOWS OF BIRATIONAL QUADRATIC TRANSFORMATIONS IN $\mathbb{P}^3$

MÒNICA MANJARÍN

ABSTRACT. We study and classify up to a linear conjugation germs of flows in the space of quadratic birational transformations of the complex projective space of dimension 3. As a consequence we show that every quadratic flow preserves a pencil of planes through a line.

1. INTRODUCTION

Let $\mathbb{P}^m$ denote the complex projective space of dimension m . Consider $m + 1$ polynomial functions $\varphi_i(x_0, \dots, x_m)$ of $m + 1$ variables with complex coefficients, homogeneous of the same degree, without common factors and not all identically zero. They define a *rational transformation* $\varphi \in \mathbf{Rat}^m$ by

$$\varphi : \mathbb{P}^m \dashrightarrow \mathbb{P}^m; \quad [x_0, \dots, x_m] \dashrightarrow [\varphi_0(x_0, \dots, x_m), \dots, \varphi_m(x_0, \dots, x_m)].$$

We define the degree of φ , denoted by $\deg(\varphi)$, as the degree of the polynomials φ_i . We say that φ is a *birational transformation* or a *Cremona transformation* if there exists a rational transformation $\psi : \mathbb{P}^m \dashrightarrow \mathbb{P}^m$ such that $\psi \circ \varphi$ is the identity. The study of these transformations, introduced by L. Cremona, was initiated in the 19th century by Cremona, Noether and Jonquières among others. The group of birational transformations of $\mathbb{P}^m$ or *Cremona group* will be denoted by $\mathbf{Bir}^m$.

We denote by $\text{Ind}(\varphi)$ the *indeterminacy locus* of φ , i.e. the points where all the polynomials $\varphi_0, \dots, \varphi_m$ vanish. We define the *Jacobian* of φ as the determinant of the matrix $(\frac{\partial \varphi_i}{\partial x_j})$ and it is denoted by $\text{Jac}(\varphi)$. Finally $\text{Exc}(\varphi)$ denotes the *exceptional locus* of φ , i.e. zero set of $\text{Jac}(\varphi)$.

Example 1.1. The *Cremona involution* σ in $\mathbb{P}^2$ is defined by $\sigma[x, y, z] = [yz, xz, xy]$. Its indeterminacy locus consists of three distinct points $P_0 = [1, 0, 0]$, $P_1 = [0, 1, 0]$ and $P_2 = [0, 0, 1]$ and σ blows down the three coordinate lines $x = 0$, $y = 0$ and $z = 0$, i.e. $\text{Ind}(\varphi) = \{P_0, P_1, P_2\}$ and $\text{Exc}(\varphi) = \{[x, y, z] \in \mathbb{P}^2 : x \cdot y \cdot z = 0\}$.

In $\mathbb{P}^2$ birational transformations have been extensively studied, for monographs on the subject we refer the reader to [14], [13], [1] and [9]. Among the most significant results we find the following classical theorem proved by Noether in 1871:

Theorem 1.2 (Noether). $\mathbf{Bir}^2$ is generated by $\text{Aut}_{\mathbb{C}}(\mathbb{P}^2) = \text{PGL}(3, \mathbb{C})$ and the Cremona involution σ .

Alternatively, Noether's theorem can be stated by saying that up to an element of $\text{PGL}(3, \mathbb{C})$ every birational transformation of $\mathbb{P}^2$ is a composition of quadratic transformations.

In sharp contrast, in $\mathbb{P}^m$ for $m \geq 3$, I. Pan proved the following in [17] (a similar result had also been previously proved by H. Hudson for $m = 3$, see [14]):

Theorem 1.3 (Pan). Let $m \geq 3$. Every set of generators of $\mathbf{Bir}^m$ must contain an uncountable number of transformations of degree > 1 .

This result gives us a hint of the reason why so little is known on Cremona transformations in higher dimensions.

There is a certain number of classical problems regarding the Cremona group, although they have been addressed mostly in dimension 2. We mention briefly some of them, the bibliography included is by no means exhaustive, see the monographs mentioned above for more references. On the one hand we find the study of special families of transformations, such as involutions (see [3] or [2]), De Jonquières maps, i.e. elements of the plane Cremona group which preserve a fixed rational fibration (see [15] for a presentation of the Cremona group with generators the De Jonquières maps and $\text{Aut}_{\mathbb{C}}(\mathbb{P}^1 \times \mathbb{P}^1)$, for instance) or transformations

Date: 18th February 2010.

Key words and phrases. birational quadratic transformations, Cremona group, germs of flows.

that preserve a fixed curve. Castelnuovo proved in 1892 (c.f. [6]) that an element of $\mathbf{Bir}^2$ preserving an irreducible curve of genus > 1 must be either conjugated to a De Jonquières map or of degree 2, 3 or 4 (see also [4] for a more recent and precise version). On the other hand, Kantor began the classification of finite subgroups up to a birational conjugation (c.f. [16]), a history of the classification of these groups can be found in the recent paper [10]. Demazure and Umemura studied in the 70's and the 80's (see [8] and [19], [20], and [21] respectively) algebraic subgroups of $\mathbf{Bir}^m$ of maximal rank (for $m = 2$ and $m = 3$ the problem goes back again to the end of the 19th century with the work of Enriques and Fano, see [11] and [12]). More recently, Cerveau-Déserti have introduced a new approach to study maximal uncountable Abelian algebraic subgroups of $\mathbf{Bir}^2$ (cf. [7]) by means of germs of flows.

Definition 1.4. A *germ of flow* in $\mathbf{Bir}^m$ is a germ of holomorphic map $t \mapsto \varphi_t \in \mathbf{Bir}^m$ such that

$$\varphi_{t+s} = \varphi_t \circ \varphi_s \quad \text{and} \quad \varphi_0 = \text{id}.$$

From now on φ_t will denote a germ of flow and we will talk of *flow* for shortness.

We denote by $\langle \varphi_t \rangle \subset \mathbf{Bir}^m$ the subgroup generated by $\{\varphi_t\}$ and by $\overline{\langle \varphi_t \rangle}^Z$ its Zariski closure in $\mathbf{Bir}^m$. Cerveau-Déserti have studied the maximal Abelian algebraic subgroup of $\mathbf{Bir}^2$ which contains $\overline{\langle \varphi_t \rangle}^Z$. Note that if G is a subgroup of $\mathbf{Bir}^m$, then its Zariski closure in $\mathbf{Bir}^m$ is algebraic and therefore has a finite number of connected components. Thus, if G is uncountable the identity component of its closure $\overline{G}^Z$ must have dimension at least 1, which implies that G contains flows.

They essentially tackle the cases where the transformations are of small degree, that is quadratic or cubic. Although birational transformations of a given degree of $\mathbf{Bir}^m$ do not form a subgroup (note that the degree of the composition might be bigger) they contain groups. In [7] they classify the quadratic flows up to linear conjugation and prove that every germ of flow (of arbitrary degree) preserves a fibration by lines. Their notation of flow (drooping the germ) is justified by the classification, that shows that the time coordinate t can actually be taken in $\mathbb{C}$.

The main tool consists roughly speaking in understanding the behavior of the families of points obtained taking the indeterminacy points of each quadratic transformation in the flow φ_t and the families of curves obtained taking the elements of the exceptional locus of each transformation φ_t . Another crucial ingredient, which allows to derive consequences for a flow of arbitrary degree, is a result by Cantat-Favre (see [5]. Theorem 1.2) on the group of birational transformations which preserve a foliation on $\mathbb{P}^2$.

In this paper we carry out a similar (although more involved) study on $\mathbb{P}^3$, namely we classify up to linear conjugation germs of flows of quadratic birational transformations in $\mathbb{P}^3$. We recall that in $\mathbb{P}^2$ it is long been known that, up to linear automorphisms (acting on the right and on the left), i.e. up to changes of coordinates in the target and in the origin space, there are only three quadratic transformations, depending on the number of points of $\text{Ind}(\varphi)$ (which might be 3, as for the Cremona involution, 2 or 1). Recently, an analogous classification was obtained for quadratic birational transformations in $\mathbb{P}^3$ by Pan-Ronga-Vust (c.f. [18]). We shall use that result as a departing point to achieve a classification of flows of quadratic birational transformations in $\mathbb{P}^3$. As in $\mathbb{P}^2$, the type of a quadratic transformations in $\mathbb{P}^3$ up to linear automorphisms is determined by its indeterminacy locus. Note that we shall not be able to make use of any result regarding foliations on $\mathbb{P}^3$, for no such result is available, and hence we will restrict ourselves to the case of degree 2.

Recall that the degree of the inverse of a birational transformation φ of $\mathbb{P}^m$ is lower or equal than $(\deg \varphi)^{m-1}$. The pair $(\deg(\varphi), \deg(\varphi^{-1}))$ is called *bidegree of a birational transformation* φ and is denoted by $\text{bideg}(\varphi)$. Note that inverses of quadratic transformations in $\mathbb{P}^2$ are always quadratic, while in $\mathbb{P}^3$ if we exclude the linear transformations we obtain three possibilities for the bidegree: $(2, 2)$, $(2, 3)$ and $(2, 4)$. We denote by $\mathbf{Bir}_2^3$ the set of quadratic birational transformations of $\mathbb{P}^3$ and by $\mathbf{lin}$ the linear transformations in $\mathbf{Bir}_2^3$, which can be seen in the following form: choose $\ell_0, \ell_1, \ell_2, \ell_3, \ell \in A_1$, where A_1 denotes the vector space of linear forms on $\mathbb{C}^4$, ℓ is not zero and $\ell_0, \ell_1, \ell_2, \ell_3$ are linearly independent, then $(\ell\ell_0, \ell\ell_1, \ell\ell_2, \ell\ell_3) \in \mathbf{Bir}_2^3$. The subset of $\mathbf{Bir}_2^3$ of transformations of bidegree $(2, 2)$, which also contains $\mathbf{lin}$, is denoted by $\mathbf{Bir}_{2,2}^3$.

Note that if φ_t is a germ of flow in $\mathbf{Bir}_2^3$ then $\varphi_t \in \mathbf{Bir}_{2,2}^3$. Indeed, given $\varphi_t \in \mathbf{Bir}_2^3$ its inverse φ_{-t} must be quadratic. It follows that for our purposes we can focus our attention on the transformations of $\mathbf{Bir}_{2,2}^3$, rather than on $\mathbf{Bir}_2^3$.

Definition 1.5. Let φ_t be a flow in $\mathbf{Bir}^3$. We say that φ_t is *quadratic* if $\varphi_t \in \mathbf{Bir}_{2,2}^3$ and it is not contained in $\mathbf{lin}$ for a generic t .

The indeterminacy locus of a Cremona transformation $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$ is either $C_I \cup P_I$ or C_I , where C_I is a plane conic and P_I a point. In the first case we say that φ is *generic* (and the point P_I is not contained in the plane of the conic) and in the second case that it is *non-generic*. Alternatively we can define P_I as the only point blown up to an hyperplane by φ_t , which determines the point P_I even in the non-generic case, in which $P_I \in C_I$. When φ_t is non-generic we distinguish two cases depending on whether the multiplicity of the conic C_I at P_I is 1 or 2. These three possibilities (generic, non-generic with multiplicity 1, and non-generic with multiplicity 2) determine three different behaviours of the germs of flows whose elements are of the corresponding types. Pan-Ronga-Vust prove that there are 7 types of transformations of $\mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$ up to changes of coordinates in the origin and target spaces that they denote by $\mathbf{gen}^{[2]}(\mathbf{O})$, $\mathbf{gen}^{[2]}(\times)$, $\mathbf{gen}^{[2]}(\//)$, $\mathbf{tan}^{[2]}(\mathbf{O})$, $\mathbf{tan}^{[2]}(\times)$, $\mathbf{tan}^{[2]}(\//)$ and $\mathbf{osc}^{[2]}(\times)$. The symbols $\mathbf{O}$, $\times$ and $\//$ make reference to the type of the conic C_I . Briefly, the first three cases correspond to the generic case, $\mathbf{tan}^{[2]}(\mathbf{O})$ and $\mathbf{tan}^{[2]}(\times)$ are non-generic with conic C_I of multiplicity 1 at P_I and $\mathbf{tan}^{[2]}(\//)$ and $\mathbf{osc}^{[2]}(\times)$ are non-generic with conic C_I of multiplicity 2 at P_I . We denote by H the plane containing C_I which is the only hyperplane contracted to a point and by S , if it exists, the only surface contracted to a curve by φ .

We devote section 2 to studying the geometrical properties of the elements of $\mathbf{Bir}_{2,2}^3$ which are necessary for our purposes, namely the indeterminacy and the exceptional loci. In section 3 we consider quadratic flows $\{\varphi_t\}$. Given a germ of flow we will consider the family $\{P_{I_t}\}$ of points, where P_{I_t} is the only point blown up to a plane by φ_t , and analogously for the families $\{C_{I_t}\}$, $\{H_t\}$ and also, when it exists, $\{S_t\}$. The behaviour of these families (whose elements belong either to the indeterminacy locus or to the exceptional locus of the corresponding φ_t) will play a central role in our proof of the classification of germs of quadratic transformations in $\mathbf{Bir}^3$. We prove the following:

Main Theorem 1. *Let φ_t be a quadratic flow in $\mathbf{Bir}^3$. Then one of the following possibilities hold:*

- a) $\varphi_t \in \mathbf{gen}^{[2]}(\//) \cup \mathbf{lin}$, P_I , S , C_I are fix and H_t is mobile,
- b) $\varphi_t \in \mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$, P_I , S are fix and H_t , C_{I_t} are mobile,
- c) $\varphi_t \in \mathbf{tan}^{[2]}(\//) \cup \mathbf{lin}$, H , C_I are fix and P_{I_t} can be either fix or mobile,
- d) $\varphi_t \in \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$, H , P_I are fix and C_I can be either fix or mobile.

Moreover, if H is fix then φ_t is a polynomial flow, i.e. $\varphi_t|_{\mathbb{P}^3 \setminus H} : \mathbb{P}^3 \setminus H \rightarrow \mathbb{P}^3 \setminus H$ is polynomial for each t . In particular there are no flows in $\mathbf{gen}^{[2]}(\mathbf{O}) \cup \mathbf{gen}^{[2]}(\times) \cup \mathbf{lin}$.

This result gives us the possible configurations and will be the first step to achieve a classification.

We also prove the following higher dimensional analogous in $\mathbb{P}^3$ of the result in $\mathbb{P}^2$ that states that any flow preserves a rational fibration (see [7]):

Main Theorem 2. *Let φ_t be a quadratic flow in $\mathbf{Bir}^3$. Then:*

- i) *There exists a line L such that φ_t preserves the family of hyperplanes through L . Moreover in cases a) and c) we can choose $L = C_I$.*
- ii) *If P_I is fix then φ_t preserves the family of hyperplanes through P_I (in particular the family of lines through P_I).*

In the generic case we deduce a geometrical description of the behaviour of a flow:

Main Corollary 3. *Let φ_t a quadratic generic flow in $\mathbf{Bir}^3$. Then φ_t is determined by a linear flow η_t on the $\mathbb{P}^2$ of the net of lines through P_I and a linear flow χ_t on the $\mathbb{P}^1$ of the pencil of planes through the line C_I . Namely*

$$\varphi_t(P) = \eta_t(P \vee P_I) \cap \chi_t(P \vee C_I).$$

Finally,

Main Theorem 4. *Let φ_t be a quadratic flow in $\mathbf{Bir}^3$. Then, up to linear conjugation, φ_t is in one (and only one) of the lists of theorems 3.20 (generic case), 3.26 (non-generic with conic C_I of multiplicity 1 at P_I) and 3.33 (non-generic with conic C_I of multiplicity 2 at P_I).*

In particular one concludes that if φ_t is a germ of quadratic flow then φ_t is defined for every $t \in \mathbb{C}$, which justifies the use of the term flow.

The author is indebted to D. Cerveau for suggesting the problem and to both him and J. Déserti for explaining their work, for helpful discussions and hints and also for comments on this manuscript. This

paper was written during a postdoctoral research stay at the IRMAR, the author would also like to thank the members of Analytic Geometry team for their warm hospitality.

2. $\mathbf{Bir}_{2,2}^3$

In this section we discuss results on $\mathbf{Bir}_{2,2}^3$, mainly from [18]. We are particularly interested in which points, curves and surfaces are contracted or exploded by a given $\varphi \in \mathbf{Bir}_{2,2}^3$. We begin by recalling an elementary result:

Lemma 2.1. *Let $S = \{s = 0\}$ be a surface contracted to a point P by $\varphi \in \mathbf{Bir}^3 \setminus \mathbf{lin}$, where s is a homogeneous polynomial. Then $\deg s = 1$, i.e. S is a hyperplane. Moreover there is at most one hyperplane contracted to a point by $\varphi \in \mathbf{Bir}_{2,2}^3$.*

Proof. Up to a change of coordinates we can assume that $P = [0, 0, 0, 1]$. Then if S is contracted to P we have

$$\varphi_0 = a_0 \cdot s, \quad \varphi_1 = a_1 \cdot s, \quad \varphi_2 = a_2 \cdot s,$$

where a_0, a_1, a_2 are homogeneous polynomials. It follows that $\deg s \leq 2$. If $\deg s = 2$ then $a_0, a_1, a_2 \in \mathbb{C}$ and $\varphi_0, \varphi_1, \varphi_2$ are linearly dependent, which contradicts the hypothesis that φ is birational. Assume now that $H = (h = 0)$ and $H' = (h' = 0)$ are two different hyperplanes contracted to a point P . As before we can assume that $P = [0, 0, 0, 1]$, then

$$\varphi_0 = a_0 \cdot h \cdot h', \quad \varphi_1 = a_1 \cdot h \cdot h', \quad \varphi_2 = a_2 \cdot h \cdot h'$$

where $a_0, a_1, a_2 \in \mathbb{C}$, so again $\varphi_0, \varphi_1, \varphi_2$ would be linearly dependent, which is a contradiction. If $\varphi(H) \neq \varphi(H')$ we would proceed in a similar way. $\square$

We recall next a result by Pan-Ronga-Vust (c.f. [18]) and we derive some consequences. We denote by A_2 the vector space of quadratic forms on $\mathbb{C}^4$. Given $\varphi \in \mathbf{Bir}_2^3$ such that in a given set of coordinates it is written as $\varphi[x_0, x_1, x_2, x_3] = [\varphi_0, \varphi_1, \varphi_1, \varphi_2]$ we define M_φ as the vector subspace of A_2 of dimension 4 generated by $(\varphi_i)_{i=1}^3$. Then classifying elements of $\mathbf{Bir}_{2,2}^3$ up to linear changes of coordinates at the origin and at the target space is equivalent to studying the action of $\mathrm{GL}(4)$ on the subspaces M of dimension 4 of A_2 such that

$$\varphi_M : \mathbb{P}^3 \dashrightarrow \mathbb{P}(M^\vee), \quad x \dashrightarrow \{f \in M : f(x) = 0\}$$

is birational, where $M^\vee$ denotes the dual space of M .

Proposition 2.2 (Pan-Ronga-Vust). *Let $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$. Then there exist a quadric $Q = \{q = 0\}$ of rank lower or equal than 3 and a plane $H = \{h = 0\}$ such that if C_I denotes the conic $Q \cap H$ then*

$$M_\varphi \subseteq \{f \in A_2 : C_I \subset (f = 0)\}.$$

Proof. Let q, q' two generic elements of M_φ . Then the intersection of the quadrics $(q = 0)$ and $(q' = 0)$ is the strict transform B by φ^{-1} of a general line and an effective 1-cycle which depends only on M and whose support is contained in $\mathrm{Ind}(\varphi)$. The curve B is rational and irreducible and as φ^{-1} is quadratic B is a smooth conic and we can assume that $B = (q(x_0, x_1, x_2) = x_3 = 0)$. Therefore there exist $h \in A_1$ and $\alpha \in \mathbb{C}$ such that $q' = \alpha q + x_3 h$. Then $B_0 = (q = h = 0)$ and $M_\varphi \subseteq \{f \in A_2 : B_0 \subset (f = 0)\}$. $\square$

Proposition 2.3. *Let $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$. With the same notation as above:*

- (i) *We can choose coordinates such that $H = (x_3 = 0)$, $Q = (q(x_0, x_1, x_2) = 0)$ and $\ell_0, \ell_1, \ell_2, \ell_3 \in A_1$ such that*

$$\varphi[x_0, x_1, x_2, x_3] = [\ell_0 x_3, \ell_1 x_3, \ell_2 x_3, q + \ell_3 x_3].$$

- (ii) *The hyperplane H is the only surface contracted to a point. We denote $P = \varphi(H)$.*
- (iii) *There is exactly one point $P_I \in \mathrm{Ind}(\varphi)$ which is blown up to a surface, which is a hyperplane and will be denoted by H_I .*
- (iv) *We denote by H^{-1} the hyperplane contracted to a point P^{-1} by φ^{-1} and by $P_I^{-1} \in \mathrm{Ind}(\varphi^{-1})$ the point blown up to a hyperplane H_I^{-1} . Then*

$$H = H_I^{-1}, \quad H_I = H^{-1}, \quad P = P_I^{-1}, \quad P_I = P^{-1}.$$

- (v) Either $\text{Ind}(\varphi) = C_I$ or $\text{Ind}(\varphi) = C_I \cup P_I$. In the second case we will say that φ is generic.
- (vi) φ is generic if and only if we can choose coordinates such that

$$\varphi[x_0, x_1, x_2, x_3] = [x_0x_3, x_1x_3, x_2x_3, q(x_0, x_1, x_2)]$$

and φ is non generic if and only if we can choose coordinates such that

$$\varphi[x_0, x_1, x_2, x_3] = [x_0x_3, x_1x_3 + q(x_0, x_1, x_2), x_2x_3, x_3^2].$$

- (vii) With the above coordinates, if φ is generic then

$$\text{Jac}(\varphi) = -2x_3^2q(x_0, x_1, x_2)$$

and if φ is non-generic then

$$\text{Jac}(\varphi) = 2x_3^3\left(x_3 + \frac{\partial q}{\partial x_1}\right).$$

- (viii) If φ is generic then the quadric $S = (q = 0)$ is contracted to a conic C . Moreover S is the only surface contracted to a curve and $q \in M_\varphi$.
- (ix) If φ is non-generic and $\frac{\partial q}{\partial x_1} \neq 0$ then the hyperplane $\Pi = (x_3 + \frac{\partial q}{\partial x_1} = 0)$ is contracted to a conic C . Moreover Π is the tangent plane to the quadric $x_1x_3 + q \in M_\varphi$ at the point $(0, 1, 0, 0) = P_I$ and it is the only surface contracted to a curve.
- (x) If φ is non-generic and $\frac{\partial q}{\partial x_1} = 0$ there exist no surfaces contracted to curves.
- (xi) The image by $\varphi \in \text{Bir}_{2,2}^3 \setminus \text{lin}$ of a hyperplane L is a hyperplane if and only if $P_I \in L$ or $L \cap H \subset C_I$ (which implies $\text{rank } C_I \leq 2$).
- (xii) If C_I is blown to a quadric $S_I \not\supset H_I$ then the image by φ of a quadric containing C_I is a quadric.
- (xiii) The strict transform by φ^{-1} of a line through P_I is a conic through P_I .

Proof. It is clear that we can choose coordinates as in (i). Using this expression we see that the hyperplane $H = (x_3 = 0)$ is contracted to the point $P = [0, 0, 0, 1]$, by the previous lemma, it must be the only surface contracted to a point, which proves (ii). To prove (iii) note that a point is blown up to a surface if and only if its inverse $\varphi^{-1} \in \text{Bir}_{2,2}^3$ contracts the surface to the point. This implies that there is exactly one point with this property, denoted P_I , and that it is blown up to an hyperplane, which we will denote by H_I . Moreover, $P_I = P^{-1}$ and $H_I = H^{-1}$ and analogously for the other equalities in (iv). We have $\varphi(H) = P$, therefore $H = \varphi^{-1}(P)$, which implies $P_I^{-1} = P$ and $H_I^{-1} = H$. From $\varphi^{-1}(H^{-1}) = P^{-1}$ we would prove the other equalities. The assertion that $\text{Ind}(\varphi)$ is either the conic C_I or the conic C_I plus a point is clear in view of the previous expression in coordinates, it is enough to prove that the extra point of $\text{Ind}(\varphi)$ in the generic case is exactly the point P_I blown up to a hyperplane. As this is an easy consequence of (ix), we proceed to prove (vi). Assume that

$$\varphi[x_0, x_1, x_2, x_3] = [\ell_0x_3, \ell_1x_3, \ell_2x_3, q + \ell_3x_3].$$

A point R belongs to $\text{Ind}(\varphi) \setminus C_I$ if it verifies the equations:

$$\ell_0x_3 = \ell_1x_3 = \ell_2x_3 = q + \ell_3x_3 = 0$$

and $x_3(R) \neq 0$. We now claim that if such an R exists then

$$x'_0 = \ell_0, \quad x'_1 = \ell_1, \quad x'_2 = \ell_2, \quad x'_3 = x_3$$

is a change of coordinates. Indeed, if ℓ_0, ℓ_1, ℓ_2 were not linearly independent then φ would not be birational. Moreover $\ell_0(R) = \ell_1(R) = \ell_2(R) = 0$. Therefore

$$\varphi[x'_0, x'_1, x'_2, x'_3] = [x'_0x'_3, x'_1x'_3, x'_2x'_3, q'(x'_0, x'_1, x'_2, x'_3)]$$

and $R = [0, 0, 0, 1]$. As $R \in \text{Ind}(\varphi)$ we conclude that q' has no quadratic terms in x'_3 so we can make a change of coordinates at the target space so that q' does not depend on x'_3 . On the other hand, φ is non generic if and only if ℓ_0, ℓ_1, ℓ_2 and x_3 are linearly independent. Indeed, for φ to be birational ℓ_0, ℓ_1 and ℓ_2 must be linearly independent and the equation $\ell_0 = \ell_1 = \ell_2 = 0$ has a single point R as solution. Then $R \in C_I$ if and only if $x_3(R) = q(R) = 0$. Up to a change of coordinates we can assume that $R = [0, 1, 0, 0]$ and that

$$\varphi[x'_0, x'_1, x'_2, x'_3] = [x'_0x'_3, \varphi'_1(x'_0, x'_1, x'_2, x'_3), x'_2x'_3, x'^2_3].$$

As $\varphi'_1(R) = 0$ the polynomial φ'_1 cannot have terms in x'^2_1 and we can also assume that the only terms in x'_3 are of the form $x'_1x'_3$. This allows us to conclude. The proof of (vi) is a computation. Then (vii)

is consequence of the expressions **(v)** and **(vi)**. For **(ix)** note only that $q(x_0, x_1, x_2)$ does not contain the monomial x_1^2 because $P_I = (0, 1, 0, 0) \in \text{Ind}(\varphi)$. The last statements are clear. $\square$

Let $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$ generic, there are three possibilities for $q(x_0, x_1, x_2)$ which correspond to rank of C_I equal to 3, 2 and 1. Then one can choose:

$$q_O = x_0^2 - x_1x_2, \quad q_\times = x_1x_2, \quad q_{//} = x_2^2.$$

One checks readily that with these choices of coordinates at the origin and the target space one has $\varphi = \varphi^{-1}$. Similar arguments can be applied in the rest of the cases, yielding the following theorem:

Theorem 2.4 (Pan-Ronga-Vust). *Let $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$. Up to a linear automorphism on the right and on the left one has 7 possibilities for φ :*

$$\begin{aligned} \mathbf{gen}^{[2]}(\alpha) &:= \langle x_0x_3, x_1x_3, x_2x_3, q_\alpha \rangle \\ \mathbf{tan}^{[2]}(\alpha) &:= \langle x_0x_3, x_2x_3, x_3^2, x_1x_3 - q_\alpha \rangle \\ \mathbf{osc}^{[2]}(\times) &:= \langle x_0x_3 - x_1x_2, x_1x_3, x_2x_3, x_3^2 \rangle \end{aligned}$$

where $\alpha = O, \times$ or $//$ with $q_O = x_0^2 - x_1x_2$, $q_\times = x_1x_2$, $q_{//} = x_2^2$.

Corollary 2.5. *Let $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$. Then φ and φ^{-1} are in the same class with respect to composition by linear automorphisms on the right and on the left.*

Corollary 2.6. *Let $\varphi \in \mathbf{Bir}_{2,2}^3 \setminus \mathbf{lin}$. Then:*

- i) *Assume φ is generic. Then C_I is blown up to a quadric S_I . Moreover, if φ^{-1} refers to the corresponding element for φ^{-1} , then*

$$S = S_I^{-1}, \quad C = C_I^{-1}.$$

- ii) *Assume φ is non generic and $\frac{\partial q}{\partial x_1} \neq 0$, i.e. $\varphi \in \mathbf{tan}^{[2]}(O) \cap \mathbf{tan}^{[2]}(\times)$. Then C_I is blown up to a plane Π_I . Moreover, if φ^{-1} refers to the corresponding element for φ^{-1} then*








$$\Pi = \Pi_I^{-1}, \quad C = C_I^{-1}.$$

Proof. Assume that φ is generic (for the other cases similar arguments apply). As φ and φ^{-1} are in the same class there is a surface S_I contracted to a conic $\tilde{C}$ by φ^{-1} . As $\tilde{C} \subset \text{Ind}(\varphi)$ we must have $\tilde{C} = C_I$. Thus, C_I is a conic blown up to a surface S . Moreover, $S = S_I^{-1}$ and $C = C_I^{-1}$. $\square$

In the figures of the next page we represent the main geometrical objects of each type of transformations. See the table in page 7 for a study of the seven cases. We will write φ instead of $\varphi[x_0, x_1, x_2, x_3]$ and analogously for φ^{-1} .

It will also be useful to have a geometric description of the elements of the *linear system* Γ_φ associated to φ , i.e. the linear system of inverse images of hyperplanes by φ . Note that if $\varphi = (\varphi_0, \varphi_1, \varphi_2, \varphi_3)$ for some choices of coordinates then $(\varphi_i = 0) = \varphi^{-1}(x_i = 0) \in \Gamma_\varphi$. Recall that we say that a quadric *osculates* at a point P along a germ of curve C if it has a contact of order 2 with C at P , i.e. if the multiplicity of intersection of the surface with C at p is 3).

Type	Definition of Γ_φ
gen ^[2] (O)	Quadrics containing a smooth conic C_I and a point $P_I \notin C_I$
gen ^[2] ($\times$)	Quadrics containing a conic C_I of rank 2 and a point $P_I \notin C_I$
gen ^[2] (//)	Cones containing a line C_I and a point $P_I \notin C_I$ and tangents to a plane S_I along C_I
tan ^[2] (O)	Quadrics containing a smooth conic C_I and tangents to a plane S at a point $P_I \in C_I$
tan ^[2] ($\times$)	Quadrics containing a conic C_I of rank 2 and tangents to a plane S at a point $P_I \in C_I$
tan ^[2] (//)	Cones containing a line C_I , tangents to a plane H along C_I and osculating at a point $P_I \in C_I$ along a curve α tangent to H at P_I
osc ^[2] ($\times$)	Quadrics containing a conic C_I of rank 2, $C_I = L_1 \cup L_2$, and osculating at the point $P_I = L_1 \cap L_2$ along a curve α tangent to the plane $H = L_1 \vee L_2$ at P_I


gen^[2](O)

gen^[2]($\times$)

gen^[2](//)

tan^[2](O)

tan^[2]($\times$)

tan^[2](//)

osc^[2]($\times$)

Type, expression and inverse	$\text{Ind}(\varphi)$	$\text{Jac}(\varphi), \text{Jac}(\varphi^{-1})$	P_I, H_I	H, S, Π, C
gen ^[2] (O) $\varphi = [x_0x_3, x_1x_3, x_2x_3, x_0^2 - x_1x_2]$ $\varphi^{-1} = \varphi$	$(x_3 = x_0^2 - x_1x_2 = 0)$ $\cup P_I$ with $P_I = [0, 0, 0, 1]$	$-2x_3^2(x_0^2 - x_1x_2),$ Idem	$P_I,$ $H_I = (x_3 = 0) = H$	$\varphi(H) = P = P_I,$ $S = (x_0^2 - x_1x_2 = 0),$ $\varphi(S) = S \cap H = C$
gen ^[2] ($\times$) $\varphi = [x_0x_3, x_1x_3, x_2x_3, x_1x_2]$ $\varphi^{-1} = \varphi$	$(x_3 = x_1x_2 = 0)$ $\cup P_I$ with $P_I = [0, 0, 0, 1]$	$-2x_3^2x_1x_2,$ Idem	$P_I,$ $H_I = (x_3 = 0) = H$	$\varphi(H) = P = P_I,$ $S = (x_1x_2 = 0) = 0,$ $\varphi(S) = S \cap H = C$
gen ^[2] ($//$) $\varphi = [x_0x_3, x_1x_3, x_2x_3, x_2^2]$ $\varphi^{-1} = \varphi$	$(x_3 = x_2^2 = 0)$ $\cup P_I$ with $P_I = [0, 0, 0, 1]$	$-2x_3^2x_2^2,$ Idem	$P_I,$ $H_I = (x_3 = 0) = H$	$\varphi(H) = P = P_I,$ $S = (x_2^2 = 0),$ $\varphi(S) = S \cap H = C$
tan ^[2] (O) $\varphi = [x_0x_3, x_1x_3 + x_1x_2 - x_0^2, x_2x_3, x_3^2]$ $\varphi^{-1} = [x_0(x_2 + x_3), x_1x_3 + x_0^2, x_2^2 + x_2x_3, x_2x_3 + x_3^2]$	$(x_3 = x_0^2 - x_1x_2 = 0)$	$2x_3^3(x_2 + x_3),$ $2(x_2 + x_3)^2x_3$	$P_I = [0, 1, 0, 0],$ $H_I = (x_2 + x_3 = 0)$	$H = (x_3 = 0), \varphi(H) = P = P_I,$ $\Pi = (x_2 + x_3 = 0),$ $\varphi(\Pi) = (x_2 + x_3 = x_0^2 - x_1x_3 = 0) = C$
tan ^[2] ($\times$) $\varphi = [x_0x_3, x_1x_3 - x_1x_2, x_2x_3, x_3^2]$ $\varphi^{-1} = [x_0(x_3 - x_2), x_1x_3, x_2x_3 - x_2^2, x_3^2 - x_2x_3]$	$(x_3 = x_1x_2 = 0)$	$2x_3^3(x_3 - x_2),$ $2(x_3 - x_2)^3x_3$	$P_I = [0, 1, 0, 0],$ $H_I = (x_3 - x_2 = 0)$	$H = (x_3 = 0), \varphi(H) = P = P_I,$ $\Pi = (x_2 - x_3 = 0),$ $\varphi(\Pi) = (x_3 - x_2 = x_1 = 0) = C$
tan ^[2] ($//$) $\varphi = [x_0x_3, x_1x_3 - x_2^2, x_2x_3, x_3^2]$ $\varphi^{-1} = [x_0x_3, x_1x_3 + x_2^2, x_2x_3, x_3^2]$	$(x_3 = x_2^2 = 0)$	$2x_3^4,$ Idem	$P_I = [0, 1, 0, 0],$ $H_I = (x_3 = 0) = H$	$\varphi(H) = P = P_I$
osc ^[2] ($\times$) $\varphi = [x_0x_3 - x_1x_2, x_1x_3, x_2x_3, x_3^2]$ $\varphi^{-1} = [x_0x_3 + x_1x_2, x_1x_3, x_2x_3, x_3^2]$	$(x_3 = x_1x_2 = 0)$	$2x_3^4,$ Idem	$P_I = [1, 0, 0, 0],$ $H_I = (x_3 = 0) = H$	$\varphi(H) = P = P_I$

Remark 2.7. In the case $\varphi \in \mathbf{tan}^{[2]}(//)$ we can take α as $(x_1x_3 - x_2^2 = x_0 = 0)$ (with the previous expression) and for $\varphi \in \mathbf{osc}^{[2]}(\times)$ we can choose $\alpha = (x_0x_3 - x_2^2 = x_1 - x_2 = 0)$. Note however that these choices are not unique, for instance for $\varphi \in \mathbf{tan}^{[2]}(//)$ we could also choose α as $(x_1x_3 - x_2^2 = x_0 + x_2 = 0)$.

Proposition 2.8. *Let φ be a quadratic rational map of $\mathbb{P}^3$. Then φ is birational of bidegree $(2, 2)$ if and only if one of the following possibilities holds:*

- i) $\text{Jac}(\varphi) = h^2 \cdot q$ with $h \in A_1$, $q \in A_2$ of rank ≤ 3 , $(h, q) = 1$, $\text{Sing}(q = 0) \not\subseteq (h = 0)$ and such that $\varphi(h = 0) = P$ is a point and $\varphi(q = 0) = C$ a plane conic of the same rank as C_I such that $P \notin C$. Then $\varphi \in \mathbf{gen}^{[2]}(\alpha)$ and $\alpha = O, \times, //$ with rank $\alpha = \text{rank } q$.
- ii) $\text{Jac}(\varphi) = h^3 \cdot l$ with $h, l \in A_1$, $(h, l) = 1$ and such that $\varphi(h = 0)$ is a point P and $\varphi(l = 0)$ is a conic C such that $P \in C$. Moreover there exists a point $P_I \in \text{Ind}(\varphi)$ such that for every $q \in \Gamma_\varphi$ the plane $\Pi = (l = 0)$ is the tangent plane of the quadric $(q = 0)$ at P_I . Then $\varphi \in \mathbf{tan}^{[2]}(O) \cup \mathbf{tan}^{[2]}(\times)$.
- iii) $\text{Jac}(\varphi) = h^4$ with $h \in A_1$ and such that $\varphi(h = 0)$ is a point P and there exists a point $P_I \in \text{Ind}(\varphi)$ such that the strict transform of a line in $\mathbb{P}^3$ by φ^{-1} is a curve by the point P_I . Then $\varphi \in \mathbf{tan}^{[2]}(//) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$.

Proof. It is enough to prove the converse implications.

- i) We can assume that $h = x_3$, $q = q(x_0, x_1, x_2)$, $C = (x_3 = q = 0)$ and $P = [0, 0, 0, 1]$. Note that $(q = x_3 = 0) \subset C_I$. If $(h = 0)$ is contracted to a point then we can assume that there exist $\ell_0, \ell_1, \ell_2 \in A_1$ such that $\varphi_0 = \ell_0x_3$, $\varphi_1 = \ell_1x_3$ and $\varphi_2 = \ell_2x_3$. On the other hand as $(q = 0)$ is contracted to the plane conic C we have $\varphi_3 = \alpha q$ with $\alpha \in \mathbb{C}^*$. Now, the condition $\text{Jac}(\varphi) = x_3^2q$ implies that $\ell_0, \ell_1, \ell_2 \in A_1(x_0, x_1, x_2)$ and that they are linearly independent. Therefore, we can choose coordinates $x'_0 = \ell_0$, $x'_1 = \ell_1$, $x'_2 = \ell_2$, $x'_3 = x_3$ such that

$$\varphi[x'_0, x'_1, x'_2, x'_3] = [x'_0x'_3, x'_1x'_3, x'_2x'_3, q'(x'_0, x'_1, x'_2)].$$

Then φ is clearly birational and generic.

- ii) Without loss of generality we can assume that $h = x_3$, $\varphi(x_3 = 0) = [0, 1, 0, 0] = P$, $l = x_2 + x_3$, $\varphi(l = 0) \subset (x_2 + x_3 = 0)$ and $P_I = P = [0, 1, 0, 0]$. As Π is the tangent plane at the point P_I we have that $\varphi_i = \alpha_i x_1(x_2 + x_3) + q_i(x_0, x_2, x_3)$, with $\alpha_i \in \mathbb{C}$ and $q_i \in A_2(x_0, x_2, x_3)$. Using that $(x_3 = 0)$ is contracted to the point P we conclude that there exist $\ell_0, \ell_2, \ell_3 \in A_1(x_0, x_2, x_3)$ such that $\varphi_0 = \ell_0x_3$, $\varphi_2 = \ell_2x_3$ and $\varphi_3 = \ell_3x_3$. On the other hand, $\text{Jac}(\varphi) = (x_2 + x_3)x_3^3$ implies that ℓ_0, ℓ_2, ℓ_3 are linearly independent. Finally, $\Pi = (x_2 + x_3 = 0)$ is contracted to a plane conic. Up to a linear change of coordinates at the origin and target space one can assume that

$$\varphi[x'_0, x'_1, x'_2, x'_3] = [x'_0x'_3, x'_1(x'_2 + x'_3) + \epsilon x'^2_0 + (1 - \epsilon)x'_0x'_2, x'_2x'_3, x'^2_3],$$

with $\epsilon = 0, 1$.

- iii) We can assume that $h = x_3$ and $P = P_I = [0, 1, 0, 0]$. Then $\varphi_0 = \ell_0x_3$, $\varphi_1 = \ell_1x_3 + q$, $\varphi_2 = \ell_2x_3$, $\varphi_3 = \ell_3x_3$ with $\ell_i \in A_1(x_0, x_1, x_2, x_3)$ for $i = 0, 1, 2, 3$ and $q \in A_2(x_0, x_1, x_2)$. On the other hand

$$(\varphi_i = 0) \cap (\varphi_j = 0) = (x_3 = q = 0) \cup (\ell_i = \ell_1x_3 + q = 0)$$

for $i = 0, 2, 3$. As the strict transform by φ^{-1} of the line $(x_1 = x_i = 0)$ is the curve $(\ell_i = \ell_1x_3 + q = 0) = C_i$ for $i = 0, 2, 3$, imposing $P_I \in C_i$ yields $\ell_i \in A_1(x_0, x_2, x_3)$ for $i = 0, 2, 3$ and that q has no term in x_1^2 . Therefore

$$\text{Jac}(\varphi) = \begin{vmatrix} \frac{\partial \ell_0}{\partial x_0}x_3 & \frac{\partial \ell_1}{\partial x_0}x_3 + \frac{\partial q}{\partial x_0} & \frac{\partial \ell_2}{\partial x_0}x_3 & \frac{\partial \ell_3}{\partial x_0}x_3 \\ 0 & \frac{\partial \ell_1}{\partial x_1}x_3 + \frac{\partial q}{\partial x_1} & 0 & 0 \\ \frac{\partial \ell_0}{\partial x_2}x_3 & \frac{\partial \ell_1}{\partial x_2}x_3 + \frac{\partial q}{\partial x_2} & \frac{\partial \ell_2}{\partial x_2}x_3 & \frac{\partial \ell_3}{\partial x_2}x_3 \\ \frac{\partial \ell_0}{\partial x_3}x_3 & \frac{\partial \ell_1}{\partial x_3}x_3 + \frac{\partial q}{\partial x_3} & \frac{\partial \ell_2}{\partial x_3}x_3 & \frac{\partial \ell_3}{\partial x_3}x_3 \end{vmatrix} = 2x_3^3 \left(\frac{\partial \ell_1}{\partial x_1}x_3 + \frac{\partial q}{\partial x_1} \right) \begin{vmatrix} \frac{\partial \ell_0}{\partial x_0} & \frac{\partial \ell_2}{\partial x_0} & \frac{\partial \ell_3}{\partial x_0} \\ \frac{\partial \ell_0}{\partial x_2} & \frac{\partial \ell_2}{\partial x_2} & \frac{\partial \ell_3}{\partial x_2} \\ \frac{\partial \ell_0}{\partial x_3} & \frac{\partial \ell_2}{\partial x_3} & \frac{\partial \ell_3}{\partial x_3} \end{vmatrix}.$$

It follows that $\frac{\partial q}{\partial x_1} \neq 0$ and that $\{\ell_i\}_{i=0}^3$ are linearly independent. Up to a linear change of coordinates on x_0, x_2 at the origin we can assume that q is either $-x_2^2$ or $-x_0x_2$ (or 0, which yields a linear transformation). Now, a change of coordinates at the target space allows to assume that $\ell_i = x_i$ for $i = 0, 1, 2, 3$. Therefore, φ is birational and $\varphi \in \mathbf{tan}^{[2]}(//) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$.

□

3. FLOWS IN $\mathbf{Bir}_{2,2}^3$

Let φ_t be a quadratic flow.

We denote by H_t the hyperplane contracted by φ_t to a point $P_t := \varphi_t(H_t)$, by C_{I_t} the plane conic contained in $\text{Ind}(\varphi_t)$ (recall that $C_{I_t} \subset H_t$) and by P_{I_t} the point blown up by φ_t to a hyperplane H_{I_t} .

We denote, if it exists, the surface $S_t \neq H_t$ contracted to a plane conic C_t , and by $S_{I_t} \neq H_t$, if it exists i.e if $\varphi_t \in \mathbf{gen}^{[2]}(\cdot) \cup \mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$, the surface to which is blown up the conic C_{I_t} . Note that for $S_t \neq H_t$ (resp. $S_{I_t} \neq H_{I_t}$) we mean that there is at least one t_0 such that $S_{t_0} \neq H_{t_0}$ (resp. $S_{I_{t_0}} \neq H_{I_{t_0}}$). Note also that there exists a surface $S_t \neq H$ contracted to a curve C_t if and only if the curve C_{I_t} is blown up to a surface $S_{I_t} \neq H_{I_t}$, for a map $\varphi \in \mathbf{Bir}_{2,2}^3$ and its inverse φ^{-1} are of the same type.

Note that the set of hyperplanes contracted to a point by φ_t , i.e. $\{H_t\}$, are a germ of analytic set in $\mathbb{P}^{3\vee}$ (the dual space of $\mathbb{P}^3$). We can therefore consider the family of hyperplanes contracted to a point, and since there is a unique H_t for each t we have an analytic germ of map

$$t \mapsto H_t \in \mathbb{P}^{3\vee}.$$

Moreover H_t is well defined for $t = 0$ for one can easily see that if $\varphi_0[x_0, x_1, x_2, x_3] = [\alpha x_0, \alpha x_1, \alpha x_2, \alpha x_3]$ then $H_t \rightarrow (\alpha = 0)$ when $t \rightarrow 0$.

Analogously, we can consider the germ of map $t \rightarrow S_t$ where S_t are the surfaces contracted to a conic by φ_t (which might be quadrics, hyperplanes or not exist) in the set of quadrics of $\mathbb{P}^3$ and the family of S_t . Similarly, we can consider the families of $P_t, P_{I_t}, H_t, S_{I_t}$ and C_{I_t} .

Definition 3.1. Let φ_t be a quadratic flow and H_t as above. We say that the family H_t is *fix* if it does not depend on t and that it is *mobile* otherwise. Analogously for the rest of the elements above.

Nevertheless, P_{I_t} and C_{I_t} might not be defined for $t = 0$ when the family is mobile, as we will see in the classification of flows that we shall obtain (for P_I see Theorem 3.33, case **I** and for C_{I_t} see Theorem 3.26, case **I**).

Regarding S_{I_t} we will see, also as a consequence of the classification, that if the family exists it is fix (see Theorem ??), in particular, it will be defined for $t = 0$.

Example 3.2. From the affine flow $(x, y, z) \mapsto (\frac{x}{1-tx}, y+t, z+t, 1)$ we obtain

$$\varphi_t[x_0, x_1, x_2, x_3] = [x_0x_3, (x_1 + tx_3)(x_3 - tx_0), (x_2 + tx_3)(x_3 - tx_0), x_3(x_3 - tx_0)]$$

with $\text{Ind}(\varphi_t) = (x_0 = x_3 = 0) \cup \{[1, 0, 0, 0] =: P_I\}$, $C_I = (x_0 = x_3 = 0)$, $P_t = \varphi_t(H_t) = P_I$, $H_t = (x_3 + tx_0 = 0)$, $S_t = (x_3 = 0)$ and $C_t = \varphi_t(S_t) = C_I$. In particular $\varphi_t \in \mathbf{gen}^{[2]}(//)$ for $t \neq 0$.

Example 3.3. Let

$$\varphi_t[x_0, x_1, x_2, x_3] = [x_0(x_3 + tx_0), e^t x_1 x_3 + (e^t - 1)x_2^2, x_2(x_3 + tx_0), (x_3 + tx_0)^2].$$

Then $\text{Ind}(\varphi_t) = (x_3 + tx_0 = e^t x_1 x_3 + (e^t - 1)x_2^2 = 0) = C_{I_t}$, $P_I = [0, 1, 0, 0]$, $H_t = (x_3 + tx_0 = 0)$ and $S_t = (x_3 = 0)$. In particular $\varphi_t \in \mathbf{tan}^{[2]}(\mathbf{O})$ for $t \neq 0$.

Example 3.4. Let

$$\varphi_t[x_0, x_1, x_2, x_3] = [x_0(x_3 + tx_0), e^t x_1 x_3, x_2(x_3 + tx_0), (x_3 + tx_0)^2].$$

Then $\text{Ind}(\varphi_t) = C_{I_t} = (x_1 = x_3 + tx_0 = 0) \cup (x_3 = x_0 = 0)$, $H_t = (x_3 + tx_0 = 0)$, $P_I = [0, 1, 0, 0]$ and $S_t = (x_3 = 0)$. In particular $\varphi_t \in \mathbf{tan}^{[2]}(\times)$ for $t \neq 0$.

Example 3.5. Let

$$\varphi_t[x_0, x_1, x_2, x_3] = [tx_3^2 + x_0x_2, (x_1 + tx_2)x_2, x_2^2, x_2x_3].$$

Then $\text{Ind}(\varphi_t) = (x_2 = x_3^2 = 0)$, $H_t = (x_2 = 0)$, $P_t = \varphi_t(H_t) = [1, 0, 0, 0] = P_I$ and $C_I = (x_2 = x_3^2 = 0)$. In particular $\varphi_t \in \mathbf{tan}^{[2]}(//)$ for $t \neq 0$.

Example 3.6. Let

$$\varphi_t[x_0, x_1, x_2, x_3] = [x_0x_3 - tx_1x_2, x_1x_3, x_2x_3, x_3^2].$$

Then $\text{Ind}(\varphi_t) = (x_3 = x_1x_2 = 0) = C_I$, $H = H_I = (x_3 = 0)$ and $P_I = [1, 0, 0, 0]$. In particular $\varphi_t \in \mathbf{osc}^{[2]}(\times)$ for $t \neq 0$.

In fact we will show that there are no flows in $\mathbf{gen}^{[2]}(\mathbf{O}) \cup \mathbf{gen}^{[2]}(\times) \cup \mathbf{lin}$.

Remark 3.7. Let φ_t be a quadratic flow. Then:

- i) P_t is fix if and only if P_{I_t} is fix. Moreover in this case $P = P_I$.
- ii) H_t is fix if and only if H_{I_t} is fix. Moreover in this case $H = H_I$.
- iii) C_t is fix if and only if C_{I_t} is fix. Moreover in this case $C = C_I$.
- iv) S_t is fix if and only if S_{I_t} is fix. Moreover in this case $S = S_I$.

Note that the condition that H_t is fix, i.e. $H_t = H$, does clearly not imply that $P_t = \varphi_t(H)$ is fix.

Proof. It is a consequence of corollary 2.3, since $P_{I_t} = P_t^{-1} = P_{-t}$. One proceeds analogously to prove the other cases. $\square$

Lemma 3.8. *Let φ_t be a quadratic flow. If P_{I_t} is a mobile point blown up by φ_t into H_{I_t} , then H_{I_t} is fix.*

Proof. We have

$$\varphi_t(\varphi_s(P_{I_t})) = \varphi_s(\varphi_t(P_{I_t})) = \varphi_s(H_{I_t}).$$

Note that as P_{I_t} is mobile we can assume that $\varphi_s(P_{I_t})$ is a point. We will discuss the different possibilities:

- 1) $\varphi_t(\varphi_s(P_{I_t}))$ is a point and $H_{I_t} = H_s$, then $H_{-t} = H_s$ and we conclude that H_t is fix.
- 2) $\varphi_s(P_{I_t}) = P_{I_t}$ and $\varphi_s(H_{I_t}) = H_{I_t}$. However if H_{I_t} is mobile then $\varphi_s(H_{I_t})$ must be generically a quadric (note that $\varphi_s(P_{I_t}) = H_{I_t}$ is a contradiction for P_{I_t} is mobile and this would imply $P_{I_t} = P_{I_s}$) unless $P_{I_s} \in H_{I_t}$ or $C_{I_s} \subset H_{I_t}$ for every t, s . All is left to do is to exclude these two cases. Note that it is equivalent to assume that $P_{I_s} \in H_t$ or that $C_{I_s} \subset H_t$ respectively for every t, s for $H_t = H_{I_{-t}}$.

- Let us begin by showing that there are no quadratic flows φ_t such that P_{I_t}, H_t are mobile, $P_{I_t} \in H_s$, $\varphi_s(P_{I_t}) = P_{I_t}$ and $\varphi_s(H_t) = H_t$.

Note that the flow φ_t must be non-generic. Moreover, we must have $P_{I_t} \subset L$ line and H_t contained in the family of hyperplanes through L . The points of the segment of L described by P_{I_t} are fixed by φ_s and the plans through L too. Without loss of generality we can assume that $L = (x_0 = x_1 = 0)$. Then $H_t = (\alpha_t x_0 + \beta_t x_1 = 0)$ with $\alpha_t, \beta_t \in \mathbb{C}$ and

$$\varphi_t[x_0, x_1, x_2, x_3] = [x_0 \cdot \ell_t, x_1 \cdot \ell_t, x_2 \cdot \ell'_t, x_3 \cdot \ell'_t]$$

with $\ell_t, \ell'_t \in A_1(x_0, x_1, x_2, x_3)$. Moreover $\text{Ind}(\varphi_t) = (x_0 = x_1 = \ell'_t = 0) \cup (x_2 = x_3 = \ell_t = 0)$. If x_0, x_1, ℓ'_t are linearly independent their intersection is a point which does not belong to $(x_2 = x_3 = \ell_t = 0)$. We conclude that

$$\ell'_t = a_t x_0 + b_t x_1; \quad \ell_t = c_t x_2 + d_t x_3.$$

Therefore this would imply that $\text{Ind}(\varphi_t)$ are two lines in $\mathbb{P}^3$ that do not intersect, which is a contradiction.

- Let now see that there are no quadratic flows φ_t such that C_I is a fix line, $C_I \subset H_t$ for every t , H_t and P_{I_t} are mobile, $\varphi_s(P_{I_t}) = P_{I_t}$ and $\varphi_s(H_t) = H_t$.

We can assume that $\varphi_t \in \mathbf{gen}^{[2]}(//) \cup \mathbf{lin}$ and that $P_{I_t} \notin C_I$ for any t , otherwise we are in the previous case. The hyperplane $S_t = C_I \vee P_{I_t}$ is contracted by φ_t into $C = C_I$. Moreover, as $\varphi_s(P_{I_t}) = P_{I_t}$ and $\varphi_s(S_t)$ must be a hyperplane through C_I (by proposition 2.3 xi) we conclude that $\varphi_s(S_t) = S_t$ for every t, s . As

$$S_{I_t} = \varphi_s(S_{I_t}) = \varphi_s \circ \varphi_t(C_I) = \varphi_t \circ \varphi_s(C_I) = \varphi_t(S_{I_s}) = S_{I_s}$$

we conclude that the hyperplanes S and S_I are fix, which contradicts $\varphi_s(S_t) = C_I$.

- 3) $\varphi_s(P_{I_t}) \subset \text{Ind}(\varphi_t) \setminus P_{I_t}$. In particular, φ_t is generic. Then $\varphi_t(\varphi_s(P_{I_t})) = P_{I_t} \vee \varphi_s(P_{I_t})$. Therefore, H_{I_t} is contracted to the line $P_{I_t} \vee \varphi_s(P_{I_t})$, which implies that $\varphi_t \in \mathbf{gen}^{[2]}(//) \cup \mathbf{lin}$ and $H_t = S_s$ for every t, s . Again we have reached a contradiction for $P_{I_t} \in S_t = H_t \not\supset P_{I_t}$.

$\square$

Corollary 3.9. *Let φ_t be a quadratic flow. If H_t is mobile then P_{I_t} is fix.*

Corollary 3.10. *Let φ_t be a quadratic flow. Then either H_t or P_{I_t} are fix.*

Lemma 3.11. *Let φ_t be a quadratic flow in $\mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$. If C_{I_t} is fix then $\text{rank } C_I \leq 1$.*

Proof. Let $\varphi_t = [\varphi_{0t}, \varphi_{1t}, \varphi_{2t}, \varphi_{3t}]$, if C_I has rank greater or equal than 2 one has $C_I = (q = \ell = 0)$ where q is a homogeneous quadratic polynomial of rank lower or equal than 3, $\ell \in A_1$, $(\ell, q) = 1$ and $\varphi_{it} = \alpha_{it}q + \ell_{it}\ell$ where $\alpha_{it} \in \mathbb{C}$, $\ell_{it} \in A_1$. As $\varphi_0 = Id$ the quadratic polynomials $\alpha_{i0}q + \ell_{i0}\ell$ must have a common factor. There are two possibilities. If $\alpha_{i0} = 0$ for $i = 0, 1, 2, 3$ then for t small enough ℓ_{it} are linearly independent, which implies that $\text{Ind}(\varphi) = (\ell = q = 0)$, i.e., φ is not generic. Otherwise, we use that

$$\alpha_{jt}\varphi_{it} - \alpha_{it}\varphi_{jt} = (\alpha_{jt}\ell_{it} - \alpha_{it}\ell_{jt}) \cdot \ell$$

and not all the coefficients $\alpha_{jt}\ell_{it} - \alpha_{it}\ell_{jt}$ can be zero (recall that $\dim\langle \ell_{it} \rangle = 3$). As $\varphi_0 = Id$ we deduce that $q = \ell \cdot \ell'$ with $\ell' \in A_1$. Indeed,

$$(\alpha_{j0}x_i - \alpha_{i0}x_j) \cdot \alpha = \alpha_{j0}\varphi_{i0} - \alpha_{i0}\varphi_{j0} = (\alpha_{j0}\ell_{i0} - \alpha_{i0}\ell_{j0}) \cdot \ell$$

with $\alpha \in A_1$ such that $\varphi_0 = (\alpha x_0, \alpha x_1, \alpha x_2, \alpha x_3)$. It follows that $\alpha = a \cdot \ell$ with $a \in \mathbb{C}^*$. Therefore, ℓ must divide q , i.e. $q = \ell \cdot \ell'$ with $\ell' \in A_1$, which is a contradiction. $\square$

Lemma 3.12. *Let φ_t be a quadratic flow in $\mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$. Then either H_t or P_{I_t} are mobile.*

Proof. We assume that $P_I = [1, 0, 0, 0]$ is fix and that $H_t = (x_0 = 0)$ is fix. Using the expressions of the previous section we can assume that

$$\varphi_t[x_0, x_1, x_2, x_3] = \left[\frac{g_{2t}(x_1, x_2, x_3) + x_0 g_{1t}(x_1, x_2, x_3)}{x_0}, \ell_{1t}, \ell_{2t}, \ell_{3t} \right],$$

where $[x_1, x_2, x_3] \mapsto [\ell_{1t}, \ell_{2t}, \ell_{3t}]$ is a plane Cremona transformation, so clearly it is impossible to have $\varphi_0 = Id$. $\square$

Lemma 3.13. *Let φ_t be a quadratic flow. If C_{I_t} is a mobile plane conic blown up by φ_t into a surface $S_{I_t} \neq H_{I_t}$, then S_{I_t} is fix.*

Proof. We have

$$\varphi_t(\varphi_s(C_{I_t})) = \varphi_s(\varphi_t(C_{I_t})) = \varphi_s(S_{I_t}).$$

If C_t is mobile we can assume that $\varphi_s(C_{I_t})$ is a mobile curve. There are two possibilities:

- 1) $\varphi_s(C_{I_t}) = C_{I_t}$ and then either C_{I_t} is a curve blown up by φ_s into a surface $\varphi_s(S_{I_t})$, which implies $C_{I_t} = C_{I_s}$ (a contradiction for we are assuming that C_{I_t} is mobile) or $S_{I_t} = S_s = S_{I-s}$ and S_{I_t} and S_t are fix.
- 2) $\varphi_t(\varphi_s(C_{I_t}))$ is a curve and then $S_{I_t} = S_s = S_{I-s}$ and S_{I_t} and S_t are fix.

$\square$

Corollary 3.14. *Let φ_t be a quadratic flow. If there exists a mobile family of surfaces S_t contracted to a curve then C_{I_t} is fix.*

Corollary 3.15. *Let φ_t be a quadratic flow in $\mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$. Then either S_t or H_t are fix.*

Proof. We have already noted that if both H_t and S_t are mobile then P_I and C_I are fix, and since $P_I \vee C_I = S_t$ we conclude that S_t is fix, which contradicts the hypothesis. $\square$

Corollary 3.16. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(\times) \cup \mathbf{tan}^{[2]}(\mathcal{O}) \cup \mathbf{lin}$. Then either S_t or H_t are fix.*

Proof. Note that H_{I_t} mobile implies C_{I_t} mobile, whereas S_{I_t} mobile implies C_I fix. $\square$

Lemma 3.17. *Let φ_t be a quadratic flow in $\mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$. Then either S_t or H_t are mobile.*

Proof. Assume now that both H_t and S_t are fix. Without loss of generality we can choose coordinates such that $H = (x_3 = 0)$ and $S = (q(x_0, x_1, x_2) = 0)$, where q is an homogeneous polynomial of degree 2. Therefore if $\varphi_t = (\varphi_{0t}, \varphi_{1t}, \varphi_{2t}, \varphi_{3t})$ we have

$$\varphi_{it} = \alpha_{it}q + \ell_{it}x_3 \quad (*)$$

with $\ell_{it} \in A_1(x_0, x_1, x_2, x_3)$ such that $\dim\langle \ell_{it} \rangle_t = 3$ for every t such that φ_t is generic and $\alpha_{it} \in \mathbb{C}$. As $\varphi_0 = Id$ we have that

$$\alpha_{i0}q + \ell_{i0}x_3 = \ell x_i$$

for $i = 0, 1, 2, 3$ and $\ell \in A_1$. If $\alpha_{i0} = 0$ for $i = 0, 1, 2, 3$ then ℓ must be a constant multiple of x_3 and ℓ_{i0} must be linearly independent. Therefore for t small enough ℓ_{it} would be linearly independent and φ_t is not

generic, which is a contradiction. We conclude that at least one $\alpha_{i0} \neq 0$, but then ℓ would be a constant multiple of x_3 and it is easy to see that the condition $(*)$ would lead to contradiction. $\square$

3.1. Generic flows.

Theorem 3.18. *Let φ_t be a quadratic flow in $\mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$. Then*

- i) H_t is mobile and P_I , S and C_I are fix.
- ii) $\varphi_t \in \mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$.
- iii) φ_t preserves the family of hyperplanes through C_I , which contains $\{H_s\}$.
- iv) φ_t preserves the family of hyperplanes through P_I , which does not contain the family of hyperplanes through C_I .

In particular, there are no quadratic flows in $\mathbf{gen}^{[2]}(\mathcal{O}) \cup \mathbf{gen}^{[2]}(\times) \cup \mathbf{lin}$.

Proof. Assume that H_t is mobile, then S_t and P_t are fix and $P = P_I$ (by proposition 2.3 iv). Moreover, all the hyperplanes through the point P_I are sent by φ_t to hyperplanes through the point P . As φ_t and φ_s commute we have

$$\varphi_t(\varphi_s(H_t)) = \varphi_s(\varphi_t(H_t)) = \varphi_s(P) = \varphi_s(P_I) = H_s.$$

As H_t is mobile $H_t \neq H_s$ and we have the following possibilities:

- $H_t \subset S_s$ (in particular $\text{rank } C_{I_t} \leq 2$) and $\varphi_s(H_t) \subset C_{I_s}$. In this case, as H_t is mobile we must have S_t mobile, which is a contradiction.
- $H_t \not\subset S_s$. Then $\varphi_s(H_t)$ is a hyperplane sent by φ_t to the hyperplane H_s . This implies that $\varphi_s(H_t)$ contains either P_I or C_{I_t} . On the other hand $\varphi_s(H_t)$ is a hyperplane if and only if $P_I \in H_t$ or $C_{I_s} \subset H_t$. The first option is impossible for we are in the generic case. If $\text{rank } C_{I_s} \geq 2$ then $C_{I_s} \subset H_t$ implies $H_s = H_t$, which is also impossible for we are assuming that H_t is mobile. Therefore $\text{rank } C_{I_t} = 1$. Finally, as $C_{I_s} \subset H_t$ for every t and s and H_t is mobile, we conclude that C_{I_t} is fix, and that $\text{rank } C_I = 1$ and clearly φ_t preserves the family of hyperplanes which contain C_I , which includes the H_t . Moreover $\varphi_t \in \mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$.

Assume now that H_t is fix. Then S_t is mobile and C_I is fix. We have seen that a generic flow with C_I fix verifies $\text{rank } C_I = 1$. Therefore $\varphi_t \in \mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$. Without loss of generality we can assume that

$$H = (x_1 = 0), \quad C_I = (x_0 = x_1 = 0), \quad S_t = ((x_0 + a(t)x_1)^2 = 0).$$

Then if $\varphi = (\varphi_0, \varphi_1, \varphi_2, \varphi_3)$ we have

$$\varphi_{it} = \ell_{it} \cdot x_1 + \alpha_{it} \cdot (x_0 + a(t)x_1)^2$$

with $\alpha_{it} \in \mathbb{C}$, $\ell_{it} \in A_1(x_0, x_1, x_2, x_3)$ for $i = 0, 1, 2, 3$. As usual we impose the condition $\varphi_0 = \text{Id}$. For φ_{00} there are two possibilities:

- $\alpha_{00} = 0$ and $\varphi_{00} = \alpha x_0 x_1$ with $\alpha \in \mathbb{C}^*$,
- $\alpha_0 \neq 0$ and $\varphi_{00} = \alpha x_0 (x_0 + a(0)x_1)$ with $\alpha \in \mathbb{C}^*$.

In both cases we conclude that $\alpha_{i0} = 0$ for $i = 1, 2, 3$ (because otherwise a term in x_0^2 would appear). It is not difficult to see that then $\alpha_{00} = 0$ and ℓ_{i0} are linearly independent, which is a contradiction with the hypothesis of φ generic.

Let us finally prove **iv**). We know that C_I , S and P_I are fix and that H_t is mobile. Without loss of generality we can assume that $P_I = [1, 0, 0, 0]$, $S = (x_3^2 = 0)$, $C_I = (x_0 = x_3 = 0)$ and $H_t = (\ell_t(x_0, x_3) = a(t)x_0 + b(t)x_3 = 0)$. Then

$$\varphi_t[x_0, x_1, x_2, x_3] = [\alpha_t \cdot x_3^2 + \ell_t \cdot \ell_{0t}, \ell_t \cdot \ell_{1t}, \ell_t \cdot \ell_{2t}, \ell_t \cdot \ell_{3t}]$$

with $\alpha_t \in \mathbb{C}^*$ and $\ell_{it} \in A_1(x_0, x_1, x_2, x_3)$ for $i = 0, 1, 2, 3$. Moreover, $(\ell_{1t} = \ell_{2t} = \ell_{3t} = 0) = [1, 0, 0, 0]$, which implies that ℓ_{it} does not depend on x_0 for $i = 1, 2, 3$. Therefore the family of hyperplanes through P_I is invariant by φ_t , i.e. if L is a hyperplane such that $P_I \in L$ then $\varphi_t(L)$ is a hyperplane such that $P_I \in L$ then $\varphi_t(L)$ is a hyperplane for every t and $P_I \in \varphi_t(L)$ for every t . It is clear that the only hyperplane L containing C_I and P_I is $(x_3 = 0)$. $\square$

Remark 3.19. In the example 3.2 we have seen that there exist quadratic flows in $\mathbf{gen}^{[2]}(\cdot) \cup \mathbf{lin}$ such that neither the hyperplanes containing C_I nor the hyperplanes containing P_I are fixed one to one.

Theorem 3.20. *Let φ_t be a generic quadratic flow. Then up to a linear conjugation φ_t is of the following types:*

- a) $\varphi_t[x_0, x_1, x_2, x_3] = [x_0x_3, (x_0t + x_3) \cdot \ell_{1t}, (x_0t + x_3) \cdot \ell_{2t}, (x_0t + x_3) \cdot x_3]$, or
- b) $\varphi_t[x_0, x_1, x_2, x_3] = [x_0x_3 + a(t)x_3^2, (a(t)x_0 + x_3) \cdot \ell_{1t}, (a(t)x_0 + x_3) \cdot \ell_{2t}, (a(t)x_0 + x_3) \cdot x_3]$,

where $a(t) = \frac{e^{\alpha t} - 1}{e^{\alpha t} + 1}$ and $\alpha \in \mathbb{C}^*$ and $\Psi_t[x_1, x_2, x_3] = [\ell_{1t}, \ell_{2t}, x_3]$ is one of the following linear flows in $\mathbb{P}^2$:

- i) $\Psi_t[x_1, x_2, x_3] = [x_1, x_2 + tx_3, x_3]$,
- ii) $\Psi_t[x_1, x_2, x_3] = [x_1e^{\beta t}, x_2 + tx_3, x_3]$,
- iii) $\Psi_t[x_1, x_2, x_3] = [x_1 + x_2(e^{\beta t} - 1) + tx_3, x_2e^{\beta t}, x_3]$,
- iv) $\Psi_t[x_1, x_2, x_3] = [x_1e^{\gamma_1 t}, x_2e^{\gamma_2 t}, x_3]$,
- v) $\Psi_t[x_1, x_2, x_3] = [(x_1 + tx_2)e^{\beta t}, x_2e^{\beta t}, x_3]$,
- vi) $\Psi_t[x_1, x_2, x_3] = [x_1 + tx_2 + \frac{t^2}{2}x_3, x_2 + tx_3, x_3]$,

with $\beta \in \mathbb{C}^*$, $\gamma_1, \gamma_2 \in \mathbb{C}$. Moreover, given a pair of flows in the previous list they are linearly conjugated if and only if they are both of type a) iv) or b) iv) with $\gamma_1 = \gamma_2$ and the conjugation switches x_1 and x_2 .

Remark 3.21. Up to a normalisation of the time coordinate t by an homotopy one can assume that α is 1 in case a) and that either $\beta = 1$ or $\gamma_1 = 1$ or $(\gamma_1, \gamma_2) = (0, 1)$ in case b).

Proof. By theorem 3.18 we know that $\varphi \in \mathbf{gen}^{[2]}(//)$ with $P = P_I, C_I$ and S fix. We can therefore assume that $P_I = [1, 0, 0, 0]$, $C_I = (x_0 = x_3 = 0)$, $S = (x_3^2 = 0)$ and $H_t = (\ell_t(x_0, x_3) = a(t)x_0 + b(t)x_3 = 0)$, where $a(t), b(t)$ are holomorphic on t . Therefore, there exist $\alpha_t \in \mathbb{C}^*$, $\ell_{0t} \in A_1(x_0, x_1, x_2, x_3)$ and $\ell_{1t}, \ell_{2t}, \ell_{3t} \in A_1(x_1, x_2, x_3)$ such that

$$\varphi_t[x_0, x_1, x_2, x_3] = [\alpha_t x_3^2 + \ell_t \cdot \ell_{0t}, \ell_t \cdot \ell_{1t}, \ell_t \cdot \ell_{2t}, \ell_t \cdot \ell_{3t}].$$

We must impose that

$$(3.1) \quad \begin{aligned} \varphi_s(\varphi_t)(x) &= [\alpha_s \cdot \ell_t^2 \cdot \ell_{3t}^2 + \ell_s(\varphi_t) \cdot \ell_{0s}(\varphi_t), \{\ell_s(\varphi_t) \cdot \ell_t \cdot \ell_{is}(\ell_{1t}, \ell_{2t}, \ell_{3t})\}_{i=1,2,3}] \\ &= [\alpha_{t+s} x_3^2 + \ell_{t+s} \cdot \ell_{0t+s}, \{\ell_{it+s} \cdot \ell_{t+s}\}_{i=1,2,3}]. \end{aligned}$$

Note that

$$(3.2) \quad \ell_s(\varphi_t) = a(s)\alpha_t x_3^2 + \ell_t(a(s)\ell_{0t} + b(s)\ell_{3t}).$$

As $\varphi_s \circ \varphi_t$ is quadratic the components of the expression 3.1 must have a common factor $\rho_{t,s}(x)$ of degree 2 (which might depend on t and s). There are two possibilities for $\rho_{t,s}$ (up to a holomorphic function depending on t and s taking values in $\mathbb{C}^*$):

- (a) $\rho_{t,s}(x) = \ell_s(\varphi_t)$,
- (b) $\rho_{t,s}(x)$ is the product of ℓ_t and a factor $q_{t,s}(x)$ of $\ell_s(\varphi_t)$.

In the first case we would have

$$\varphi_s(\varphi_t)(x) = \left[\alpha_s \frac{\ell_t^2 \cdot \ell_{3t}^2}{\ell_s(\varphi_t)}, \{\ell_t \cdot \ell_{is}(\ell_{1t}, \ell_{2t}, \ell_{3t})\}_{i=1,2,3} \right].$$

It follows that ℓ_{t+s} is a multiple of ℓ_t by an element of $\mathbb{C}^*$, which implies that H_t is fix and contradicts the hypothesis. Therefore, we can assume that $\rho_{t,s}(x)$ is a product of ℓ_t and a factor $q_{t,s}(x)$ of $\ell_s(\varphi_t)$ and

$$(3.3) \quad \varphi_s(\varphi_t)(x) = \left[\alpha_s \cdot \ell_t \cdot \ell_{3t}^2 + \frac{\ell_s(\varphi_t) \cdot \ell_{0s}(\varphi_t)}{\ell_t}, \{\ell_s(\varphi_t) \cdot \ell_{is}(\ell_{1t}, \ell_{2t}, \ell_{3t})\}_{i=1,2,3} \right].$$

As $q_{t,s}(x)$ must be a common factor of $\ell_s(\varphi_t)$ and $\ell_t \cdot \ell_{3t}^2$ we conclude that either ℓ_t or ℓ_{3t} are factors of $\ell_s(\varphi_t)$. From 3.2 we derive that ℓ_t cannot be a factor of $\ell_s(\varphi_t)$ unless $a(t) = 0$ (which contradicts H_t mobile) or $\alpha_t = 0$ (which would imply φ_t linear). Thus ℓ_t is a factor of $\ell_{0s}(\varphi_t)$ (see expression 3.3) and ℓ_{3t} is a factor of $\ell_s(\varphi_t)$. Expression 3.2 implies that if ℓ_{3t} is a factor of $\ell_s(\varphi_t)$ then it is also a factor of $\alpha_t x_3^2 + \ell_t \cdot \ell_{0t}$. If

$$\ell_{0t} = A_0(t)x_0 + A_1(t)x_1 + A_2(t)x_2 + A_3(t)x_3$$

then

$$\ell_{0s}(\varphi_t) = A_0(s)(\alpha_t x_3^2 + \ell_t \cdot \ell_{0t}) + A_1(s)\ell_t \cdot \ell_{1t} + A_2(s)\ell_t \cdot \ell_{2t} + A_3(s)\ell_t \cdot \ell_{3t}.$$

From the fact that ℓ_t is a factor of $\ell_{0s}(\varphi_t)$ we conclude that $A_0(t) = 0$. We impose next that the quadric $\alpha_t x_3^2 + \ell_t \cdot \ell_{0t}$ on x_1, x_2, x_3 has rank 2 (for it admits ℓ_{3t} as a factor). Imposing this condition and using its expression in terms of $a(t)$, $b(t)$, α_t and $A_i(t)$ one concludes that $A_1(t) = A_2(t) = 0$. Thus $\ell_{0t} = \beta_t x_3$ and

$$(3.4) \quad \alpha_t x_3^2 + \ell_t \cdot \ell_{0t} = x_3 \cdot ((\alpha_t + \beta_t \cdot b(t))x_3 + a(t) \cdot \beta_t x_0)$$

As $\ell_{3t} \in A_1(x_1, x_2, x_3)$ is a factor of $\alpha_t x_3^2 + \ell_t \cdot \ell_{0t}$ we have $\ell_{3t} = \delta_t x_3$. Note that in particular in the first component of φ_t appear only the monomials $x_0 x_3$ and x_3^2 . From now on we will use a more natural notation. Namely,

$$\varphi_t[x_0, x_1, x_2, x_3] = [A(t)x_0 x_3 + B(t)x_3^2, \ell_t \cdot \ell_{1t}, \ell_t \cdot \ell_{2t}, \ell_t \cdot \delta_t x_3]$$

with $A(t)$, $B(t)$ depending holomorphically on t . From $\varphi_0 = Id$ we conclude that

$$A(0) \neq 0, \quad B(0) = 0, \quad a(0) = 0, \quad b(0) \neq 0$$

and a normalization allows us to assume that $A(0) = b(0) = 1$. Moreover, for small values of t one has $A(t) \neq 0$, $b(t) \neq 0$. Dividing all the components of φ_t by $A(t)$ and modifying ℓ_{1t} , ℓ_{2t} and δ_t if necessary we can assume that $A(t) = b(t) = 1$ for all t so

$$(3.5) \quad \varphi_t[x_0, x_1, x_2, x_3] = [x_0 x_3 + B(t)x_3^2, (a(t)x_0 + x_3)\ell_{1t}, (a(t)x_0 + x_3)\ell_{2t}, (a(t)x_0 + x_3)\delta_t x_3]$$

and $B(0) = a(0) = 1$. Now,

$$\ell_s(\varphi_t) = (a(s) + \delta_t a(t))x_0 x_3 + (\delta_t + a(s)B(t))x_3^2$$

and

$$(3.6) \quad \varphi_s(\varphi_t)(x) = \left[(\delta_t + a(t)B(s)\delta_t^2)x_0 x_3 + (\delta_t B(t) + \delta_t^2 B(s))x_3^2, \right. \\ \left. \frac{\ell_s(\varphi_t)}{x_3} \ell_{1s}(\ell_{it}), \frac{\ell_s(\varphi_t)}{x_3} \ell_{2s}(\ell_{it}), \frac{\ell_s(\varphi_t)}{x_3} \delta_t \delta_s x_3 \right].$$

As $\varphi_s(\varphi_t)(x) = \varphi_{t+s}(x)$ we have

$$B(t+s) = \frac{B(t) + \delta_t B(s)}{1 + \delta_t a(t)B(s)} \\ a(t+s) = \frac{a(s) + \delta_t a(t)}{\delta_t + a(s)B(t)} \\ \delta_{t+s} = \delta_s \frac{\delta_t + a(s)B(t)}{1 + a(t)B(s)\delta_t}$$

and setting

$$f(t, s) = \delta_t + a(t)B(s)\delta_t^2 \\ g(t, s) = \delta_t + a(s)B(t)$$

we have

$$\varphi_s(\varphi_t)(x) = [f(t, s)(x_0 x_3 + B(t+s)x_3^2), g(t, s)\ell_{t+s}\ell_{1s}(\{\ell_i\}_t), g(t, s)\ell_{t+s}\ell_{2s}(\{\ell_i\}_t), g(t, s)\ell_{t+s}\delta_{t+s}x_3].$$

In particular $\Psi_t[x_1, x_2, x_3] = [\ell_{1t}(x_1, x_2, x_3), \ell_{2t}(x_1, x_2, x_3), \delta_t x_3]$ is a linear flow. Using the classification of linear flows in $\mathbb{P}^2$ (cf. [7]) one concludes that

$$\delta_t = e^{\delta t}$$

and that Ψ_t belongs to the list given in the statement of the theorem. Therefore,

$$\frac{\ell_s(\varphi_t)}{x_3} \delta_t \delta_s x_3 = g(t, s)\ell_{t+s}\delta_{t+s}x_3$$

and it follows that $f(t, s) = g(t, s)$. This equality is equivalent to $a(t)B(s)\delta_t^2 = a(s)B(t)$. Thus,

$$\frac{a(t)}{B(t)}\delta_t^2 = \frac{a(s)}{B(s)},$$

which implies that $a(t)/B(t)$ is constant and that $\delta_t = 1$. Let $B(t) = c \cdot a(t)$ with $c \in \mathbb{C}$. We are left to solve the equation

$$(3.7) \quad a(t+s) = \frac{a(t) + a(s)}{1 + c \cdot a(t) \cdot a(s)}.$$

Assuming that s is a constant and differentiating with respect to t one obtains

$$\frac{\partial}{\partial t} a(t, s) = a'(t) \frac{1 - c \cdot a^2(s)}{1 + c \cdot a(t) \cdot a(s)}.$$

Simmetry on t and s implies that

$$a'(t) \frac{1 - c \cdot a^2(s)}{1 + c \cdot a(t) \cdot a(s)} = a'(s) \frac{1 - c \cdot a^2(t)}{1 + c \cdot a(t) \cdot a(s)},$$

so

$$\frac{a'(s)}{1 - c \cdot a^2(s)} = \frac{a'(t)}{1 - c \cdot a^2(t)}.$$

Again it follows that there is a constant $D \in \mathbb{C}$ such that

$$\frac{a'(t)}{1 - c \cdot a^2(t)} = D.$$

If $c = 0$ then as $a(0) = 0$ one has $a(t) = Dt$. If $c \neq 0$ then one obtains

$$a(t) = \tilde{c}^{-1} \frac{e^{\tilde{D}t} - 1}{e^{\tilde{D}t} + 1}, \quad B(t) = \tilde{c} \frac{e^{\tilde{D}t} - 1}{e^{\tilde{D}t} + 1},$$

where $\tilde{c}, \tilde{D} \in \mathbb{C}$ and $\tilde{c}^2 = c$.

Therefore either $a(t) = at$ and $B(t) = 0$ or $A(t) = a \frac{e^{\alpha t} - 1}{e^{\alpha t} + 1}$. In both cases $a \neq 0$ for otherwise ℓ_t would be fix and we can assume $\alpha \neq 0$ for the same reason.

Note that in case **a)** up to a good choice of ρ_0^0 we can assume that $a = 1$. Moreover if φ_t and $\tilde{\varphi}_t$ were conjugated we would have (it suffices to write the condition $A \circ \tilde{\varphi}_t = \varphi_t \circ A$ for the first coordinate):

$$\rho_0^0 x_0 x_3 + \rho_3^0 (t x_0 + x_3) x_3 = (\rho_0^0 x_0 + \rho_3^0 x_3) \rho_3^3 x_3$$

so $\rho_3^0 = 0$ and $\rho_3^3 = 1$.

To end the proof we must just show that given two flows of the previous list with different choices of the parameters they cannot be conjugated unless both flows are in case **a) iv)** or **b) iv)** and the conjugation switches x_1 and x_2 or unless we are in case **b)** with $\alpha = \tilde{\alpha}$ and $A \neq \tilde{A}$ (which in particular shows that we can assume $a = 1$).

Let φ_t and $\tilde{\varphi}_t$ be two flows of the previous list and $A = [\mu_0, \mu_1, \mu_2, \mu_3]$ with $\mu_i \in A_1(x_0, x_1, x_2, x_3)$ for $i = 0, 1, 2, 3$ a linear conjugation such that

$$A \circ \tilde{\varphi}_t = \varphi_t \circ A.$$

As $P_I = [1, 0, 0, 0]$, $C_I = (x_0 = x_3 = 0)$ and $S = (x_3^2 = 0)$ are fix and common for both flows we conclude that $\mu_1(x), \mu_2(x) \in A_1(x_1, x_2, x_3)$ and $\mu_3 = \mu_3^3 x_3$. Moreover as $\ell_t = (a(t)x_0 + x_3 = 0)$ is conjugated to $\tilde{\ell}_t = (\tilde{a}(t)x_0 + x_3 = 0)$ one has $\mu_0(x) = \mu_0^0 x_0 + \mu_3^0 x_3$. In particular if we define $\tilde{A}(x_1, x_2, x_3) := [\mu_1, \mu_2, \mu_3]$ we obtain a conjugation of the correponding flows Ψ_t and $\tilde{\Psi}_t$ in $\mathbb{P}^2$, which implies that $\Psi_t = \tilde{\Psi}_t$ excepting in the case **iv)** with a switch of the coordinates x_1 and x_2 .

Therefore in case **b)** one has (again writing the condition $A \circ \tilde{\varphi}_t = \varphi_t \circ A$ for the first coordinate)

$$\mu_0^0 (x_0 x_3 + \tilde{a}(t) x_3^2) + \mu_3^0 (\tilde{a}(t) x_0 + x_3) x_3 = (\mu_0^0 x_0 + \mu_3^0 x_3) \mu_3^3 x_3 + a(t) (\mu_3^3)^2 x_3^2.$$

Equivalently,

$$\mu_0^0 \mu_3^3 x_0 x_3 + (\mu_3^0 \mu_3^3 + a(t) (\mu_3^3)^2) x_3^2 = (\mu_0^0 + \tilde{a}(t) \mu_3^0) x_0 x_3 + (\tilde{a}(t) \mu_0^0 + \mu_3^0) x_3^2$$

which yields $\mu_3^0 = 0$ so

$$\mu_0^0 \mu_3^3 x_0 x_3 + a(t) (\mu_3^3)^2 x_3^2 = \mu_0^0 x_0 x_3 + \tilde{a}(t) \mu_0^0 x_3^2.$$

Using that $a(t) = a \frac{e^{\alpha t} - 1}{e^{\alpha t} + 1}$ and $\tilde{a}(t) = \tilde{a} \frac{e^{\tilde{\alpha} t} - 1}{e^{\tilde{\alpha} t} + 1}$ we conclude that $\mu_3^3 = 1$ and $\tilde{a}(t) = \mu_0^0 a(t)$ (with the exception already mentioned). $\square$

Corollary 3.22. *Let φ_t a quadratic generic flow. Then φ_t is determined by a linear flow η_t on the $\mathbb{P}^2$ of the net of lines through P_I and by a linear flow χ_t on the $\mathbb{P}^1$ of the pencil of planes through the line C_I . Namely, for a point $P \in \mathbb{P}^3$ we have*

$$\varphi_t(P) = \eta_t(P \vee P_I) \cap \chi_t(P \vee C_I).$$

Proof. With the coordinates of the previous theorem we have $P_I = [1, 0, 0, 0]$ and $C_I = (x_0 = x_3 = 0)$. If L is a line through P_I we define η_t by

$$\eta_t(L) = P_I \vee \Psi_t(L \cap (x_0 = 0)).$$

As a plane π through C_I is given by an equation $\rho x_0 + \sigma x_3 = 0$ we check easily that the strict transformation of π by φ_t is

$$\begin{aligned} (\rho - t\sigma)x_0 + \sigma x_3 &= 0 & \text{for } \alpha &= 0 \\ (\rho + a(-t)\sigma)x_0 + (a(-t)\rho + \sigma)x_3 &= 0 & \text{for } \alpha &\in \mathbb{C}^* \end{aligned}$$

with $a(t) = \frac{e^{\alpha t} - 1}{e^{\alpha t} + 1}$. Thus we define

$$\mu_t[\rho, \sigma] := \begin{cases} [\rho - t\sigma, \sigma] & \text{for } \alpha = 0 \\ [\rho + a(-t)\sigma, a(-t)\rho + \sigma] & \text{for } \alpha \in \mathbb{C}^* \end{cases}.$$

□

3.2. Flows in $\mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$.

Lemma 3.23. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$ or $\mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{lin}$. Then either S_t or H_t are mobile.*

Proof. Assume that both S_t and H_t are fix, then $L = H \cap S$ is a fix line and $P_{I_t} \in L$. Without loss of generality we can choose coordinates such that $H = (x_3 = 0)$ and $S = (x_0 = 0)$, then $P_{I_t} = [0, 1, f(t), 0]$. Therefore

$$\varphi_t[x_0, x_1, x_2, x_3] =$$

$$[x_3 \cdot \ell_{0t}, a_t x_0 x_1 + r_t(x_0, x_3, f(t)x_1 - x_2), b_t x_0 x_1 + s_t(x_0, x_3, f(t)x_1 - x_2), x_3 \cdot \ell_{3t}]$$

where $r_t, s_t \in A_2(x_0, x_3, f(t)x_1 - x_2)$ (recall that $\varphi_t(H) = P_{I-t} = P_t$ and that the tangent plane of $(\varphi_{it} = 0)$ at P_{I_t} is $x_0 = 0$). Then $\varphi_0 = \text{Id}$ implies that $x_3 \cdot \ell_{00} = x_0 \cdot \ell$ with $\ell \in A_1(x_0, x_1, x_2, x_3)$. Therefore $\ell = \alpha \cdot x_3$ with $\alpha \in \mathbb{C}^*$. Now it is not difficult to reach a contradiction with

$$a_0 x_0 x_1 + r_0(x_0, x_3, f(t)x_1 - x_2) = \alpha x_1 x_3.$$

□

Theorem 3.24. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{lin}$ or $\mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$. Then:*

- i) H_t and C_{I_t} are mobile and P_I and S_I are fix.
- ii) $L_{I_t} = S \cap H_t$ is a fix line through P_I .
- iii) φ_t preserves the family of hyperplanes through L_I , which includes $\{H_s\}$.
- iv) φ_t preserves the family of hyperplanes through P_I , which includes the family of hyperplanes through L_I .

Remark 3.25. In this case as P_I is contained in the line L_I we cannot determine φ_t using flows on the net of lines through P_I and the pencil of planes through L_I as in Corollary 3.22. The list in Theorem 3.33 shows that there are not other choices of a pair point/line which allows to do so either.

Proof. By corollary 3.16 and lemma 3.23 to show i) it is enough to prove that H_t is mobile and S fix. Note that if H_t mobile and S_I fix it is clear that C_{I_t} must be mobile. Assume that H is fix and S_{I_t} mobile. Then C_I is fix and

$$S_{I_{t+s}} = \varphi_{t+s}(C_I) = \varphi_t \circ \varphi_s(C_I) = \varphi_t(S_{I_s}).$$

As S_{I_t} are hyperplanes we conclude that either $P_{I_t} \in S_{I_s}$ for every t, s or $C_I \subset S_{I_t}$ for every t .

As C_I is a fix plane conic of rank greater or equal than 2 the second possibility contradicts S_t mobile unless $C_I = L_1 \cup L_2$, L_1, L_2 lines and $L_1 \subset S_{I_t}$ for every t .

If $P_{I_t} \in S_{I_s}$ for every t, s and rank $C_I = 3$ it is not difficult to conclude that P_I must be fix, which contradicts H_I fix.

If $P_{I_t} \in S_{I_s}$ for every t, s and rank $C_I = 2$, i.e. $C_I = L_1 \cup L_2$, we conclude that $P_{I_t} \in L_1$ and $L_1 \subset S_t$ for every t .

Assume then that $C_t = L_1 \cup L_2$, L_1, L_2 lines and $L_1 \subset S_t$ for every t . Without loss of generality we can choose coordinates such that

$$H = (x_3 = 0), \quad L_1 = (x_1 = x_3 = 0), \quad L_2 = (x_2 = x_3 = 0).$$

Then $P_{I_t} = [f(t), 0, 1, 0]$ and $S_t = (a(t)x_1 + b(t)x_3 = 0)$, where $a(t), b(t)$ are germs of holomorphic functions on t . If $\varphi_t = (\varphi_{0t}, \varphi_{1t}, \varphi_{2t}, \varphi_{3t})$ using $\text{Ind}(\varphi_t) = L_1 \cup L_2$ and that $(\varphi_{it} = 0)$ are quadrics tangent to S_t at the point P_{I_t} we obtain

$$\begin{aligned} \varphi_t[x_0, x_1, x_2, x_3] = \\ [\alpha_{0t} \cdot x_2(a(t)x_1 + b(t)x_3) + x_3 \cdot \ell_{0t}, x_3 \cdot \ell_{1t}, \alpha_{2t} \cdot x_2(a(t)x_1 + b(t)x_3) + x_3 \cdot \ell_{2t}, x_3 \cdot \ell_{3t}] \end{aligned}$$

with $\alpha_{it} \in \mathbb{C}$ and $\ell_{it} \in A_1(x_1, x_3, x_0 - f(t)x_2)$.

The condition $\varphi_0 = \text{Id}$ implies that (dividing by an element of $\mathbb{C}^*$ if necessary)

$$\varphi_0 = [x_0x_3, x_1x_3, x_2x_3, x_3^2]$$

so

$$\begin{aligned} x_0x_3 &= \alpha_{00}x_2(a(0)x_1 + b(0)x_3) + x_3\ell_{00} \\ x_1 &= \ell_{10} \\ x_2x_3 &= \alpha_{20}x_2(a(0)x_1 + b(0)x_3) + x_3\ell_{20} \\ x_3 &= \ell_{30} \end{aligned}$$

and $\ell_{i0} \in A_1(x_0 - f(0)x_2)$ for $i = 1, 2, 3, 4$. One concludes that

$$\ell_{00} = \beta_0(x_0 - f(0)x_2), \quad a(0) = 0, \quad \alpha_{00}b(0) - \beta_0f(0) = 0, \quad \ell_{20} = 0, \quad \ell_{10} = x_1, \quad \ell_{30} = x_3, \quad \alpha_{00}b(0) = 1.$$

We can therefore assume that $\ell_t = a(t)x_1 + x_3$ and the previous conditions for $t = 0$ translate into

$$\alpha_{00} = f(0), \quad \ell_{00} = x_0 - f(0)x_2, \quad \ell_{10} = x_1, \quad \alpha_{20} = 1, \quad \ell_{20} = 1, \quad \ell_{30} = x_3.$$

On the other hand $\varphi_t(H) = \varphi_t(x_3 = 0) = P_t = P_{I-t} = [f(-t), 0, 1, 0]$, which implies that

$$\alpha_{0t} = f(-t)\alpha_{2t}.$$

Normalizing if necessary we have

$$(3.8) \quad \varphi_t(x) = [(f(-t)x_2\ell_t + x_3\ell_{0t}, x_3\ell_{1t}, x_2\ell_t + x_3\ell_{2t}, x_3\ell_{3t})]$$

and

$$(3.9) \quad \ell_{00} = x_0 - x_2, \quad \ell_{10} = x_1, \quad \ell_{20} = 0, \quad \ell_{30} = x_3, \quad f(0) = 1.$$

Therefore

$$(3.10) \quad \begin{aligned} \varphi_s(\varphi_t(x)) = [f(-s)(x_2\ell_t + x_3\ell_{2t})\ell_s(\varphi_t) + x_3\ell_{3t}\ell_{0s}(\varphi_t), \\ x_3\ell_{3t}\ell_{1s}(\varphi_t), (x_2\ell_t + x_3\ell_{2t})\ell_s(\varphi_t) + x_3\ell_{3t}\ell_{2s}(\varphi_t), x_3\ell_{3t}\ell_{3s}(\varphi_t)] \end{aligned}$$

and

$$(3.11) \quad \ell_s \circ \varphi_t(x) = x_3(a(s)\ell_{1t} + \ell_{3t}).$$

Dividing by x_3 we obtain

$$(3.12) \quad \begin{aligned} \varphi_s \circ \varphi_t(x) = \left[f(-s)(x_2\ell_t + x_3\ell_{2t}) \frac{\ell_s(\varphi_t)}{x_3} + \ell_{3t}\ell_{0s}(\varphi_t), \right. \\ \left. \ell_{3t}\ell_{1s}(\varphi_t), (x_2\ell_t + x_3\ell_{2t}) \frac{\ell_s(\varphi_t)}{x_3} + \ell_{3t}\ell_{2s}(\varphi_t), \ell_{3t}\ell_{3s}(\varphi_t) \right]. \end{aligned}$$

As $\varphi_s \circ \varphi_t = \varphi_{t+s}$, which is a quadratic flow, we must have a linear form dividing all the components of $\varphi_s(\varphi_t)$. This form can be either **(1)** ℓ_{3t} or **(2)** another linear form $\tilde{\ell}_t$.

Let us study case **(1)**. We see that ℓ_{3t} divides either $\frac{\ell_s(\varphi_t)}{x_3} = a(s)\ell_{1t} + \ell_{3t}$ or $x_2\ell_t + x_3\ell_{2t}$. We can clearly exclude the first possibility. On the other hand

$$x_2\ell_t + x_3\ell_{2t} = x_2(a(t)x_1 + x_3) + x_3\ell_{2t}(x_1, x_3, x_0 - f(t)x_2)$$

and taking into account that $\ell_{3t} \in A_1(x_1, x_3, x_0 - f(t)x_2)$ one can check that ℓ_{3t} cannot divide this term either.

Assume now that we are in case **(2)**, i.e. there exists a form $\tilde{\ell}_t$ which is not a multiple of ℓ_{3t} which divides all the components of $\varphi_s \circ \varphi_t$. In particular $\tilde{\ell}_t$ must divide $\ell_{1s}(\varphi_t)$, $\ell_{3s}(\varphi_t)$ and $x_3(\ell_{0s}(\varphi_t) - f(-s)\ell_{2s}(\varphi_t))$. Note that if $\tilde{\ell}_t = x_3$ then $\ell_{1t}, \ell_{3t} \in A_1(x_1, x_3)$. But in this case $\frac{\ell_{3s}(\varphi_t)}{x_3} \in A_1(x_1, x_3)$ and we easily reach a contradiction. Therefore $\tilde{\ell}_t$ divides also $\ell_{0s}(\varphi_t) - f(-s)\ell_{2s}(\varphi_t)$.

If $\ell_{1t} = A_1^0(t)(x_0 - f(t)x_2) + A_1^1(t)x_1 + A_1^3(t)x_3$ then

$$\ell_{1s}(\varphi_t) = A_1^0(s)((f(-t) - f(s))x_2\ell_t + x_3(\ell_{0t} - f(s)\ell_{2t})) + A_1^1(s)x_3\ell_{1t} + A_1^3(s)x_3\ell_{3t}.$$

A direct computation shows that for $\ell_{1s}(\varphi_t)$ to have rank 2 either

$$(3.13) \quad A_1^0(s)(f(-t) - f(s))a(t) = 0 \quad \text{and} \quad \tilde{\ell}_t = x_3$$

or

$$(3.14) \quad A_1^0(s)((f(-t) - f(s))x_2 + \ell_{0t} - f(s)\ell_{2t}) + A_1^1(s)\ell_{1t} + A_1^3(s)\ell_{3t} = 0 \quad \text{and} \quad \tilde{\ell}_t = x_1 \text{ or } x_2.$$

An analogous argument applies to $\ell_{3s}(\varphi_t)$ and $\ell_{0s}(\varphi_t) - f(-s)\ell_{2s}(\varphi_t)$. As we have already discussed the possibility $\tilde{\ell}_t = x_3$ we focus on 3.14. The linear forms ℓ_{1t} , ℓ_{3t} and $\ell_{0t} - f(-t)\ell_{2t}$ are linearly independent because

$$\text{Ind}(\varphi_t) = \{\ell_{1t} = \ell_{3t} = \ell_{0t} - f(-t)\ell_{2t} = x_2\ell_t + x_3\ell_{2t} = 0\}$$

and as φ_t is not generic $\text{Ind}(\varphi_t) = L_1 \cup L_2$ for every $t \neq 0$. Now, on 3.14 we take $s = -t$. Then

$$A_1^0(-t)(\ell_{0t} - f(-t)\ell_{2t}) + A_1^1(-t)\ell_{1t} + A_1^3(-t)\ell_{3t} = 0$$

and as ℓ_{1t} , ℓ_{3t} and $\ell_{0t} - f(-t)\ell_{2t}$ are linearly independent we conclude that

$$A_1^0(-t) = A_1^1(-t) = A_1^3(-t) = 0$$

so $\ell_{1t} = 0$, which is a contradiction. We have thus proved that H_t is mobile and that P and S are fix.

Therefore we can assume that H_t and C_{It} are mobile and $S = S_I$, $P = P_I$ are fix. Let us now show that the line $L_{It} := S_I \cap H_t$ is fix. As L_{It} is a line by the point P_I contained in H_t we have $\varphi_t(L_{It}) = P_I$. Thus,

$$H_I = \varphi_t(P_I) = \varphi_s(\varphi_t(L_{Is})) = \varphi_t(\varphi_s(L_{It})).$$

It follows that $\varphi_t(L_{Is}) = P_I$, which implies that $L_{Is} \subset H_t$ for any t, s . As H_t is mobile we have L_I fix.

We are left to prove statements **iii)** and **iv)**. We can assume that $P = [0, 1, 0, 0] = P_I$, $L_I = (x_0 = x_3 = 0)$, $S = (x_3 = 0)$ and $H_t = (\ell_t = a(t)x_0 + b(t)x_3 = 0)$, where $a(t)$ and $b(t)$ are germs of holomorphic functions. Then

$$\varphi_t[x_0, x_1, x_2, x_3] = [\ell_{0t} \cdot \ell_t, \alpha_t x_1 x_3 + q_t(x_0, x_2, x_3), \ell_{2t} \cdot \ell_t, \ell_{3t} \cdot \ell_t]$$

with $\ell_{it} \in A_1(x_0, x_2, x_3)$ (recall that all the elements of the linear system Γ_{φ_t} are quadrics tangent to the plane S at the point P). In particular the statement in **iv)** is proved.

As $\varphi_0 = \text{Id}$ we conclude (dividing by an element of $\mathbb{C}^*$ if necessary) that

$$\alpha_0 = 1, q_0 = 0, \ell_{00} = x_0, \ell_{20} = x_2, \ell_{30} = x_3, \ell_{10} = x_3, a(0) = 0, b(0) = 1.$$

We can therefore assume that $b(t) = 1$. Moreover,

$$(3.15) \quad \begin{aligned} \varphi_s(\varphi_t(x)) &= [\ell_t \cdot \ell_{0s}(\ell_{it})\ell_s(\ell_{it}), \\ &\quad \alpha_s \alpha_t x_1 x_3 \ell_{3t} + \alpha_s q_t(x_0, x_2, x_3) \ell_{3t} + \ell_t q_s(\ell_{it}), \ell_t \ell_{2s}(\ell_{it})\ell_s(\ell_{it}), \ell_t \ell_{3s}(\ell_{it})\ell_s(\ell_{it})] \end{aligned}$$

where $\ell_t = a(t)x_0 + x_3$ and

$$\ell_s(\ell_{it}) = a(t)\ell_{0t} + \ell_{3t}.$$

The components of $\varphi_s \circ \varphi_t$ must admit a common linear factor for φ_{t+s} is quadratic and the factor must be either ℓ_t or $\ell_s(\ell_{it})$ (up to an element of $\mathbb{C}^*$). On the other hand from the coefficient of $x_1 x_3$ one concludes that the factor must be ℓ_{3t} . Therefore, the possibilities are:

- (I) $\ell_{3t} = \beta_t \cdot \ell_t$,
- (II) $\ell_{3t} = \beta_t \cdot \ell_s(\ell_{it})$.

The second one would imply that H_t is fix so we can assume that **(I)** holds. Therefore,

$$\varphi_s(\varphi_t(x)) = [\ell_{0s}(\ell_{it})\ell_s(\ell_{it}), \alpha_s\alpha_t\beta_tx_1x_3 + \beta_t\alpha_sq_t(x_0, x_2, x_3) + q_s(\ell_{it}), \ell_{2s}(\ell_{it})\ell_s(\ell_{it}), \beta_t\ell_s^2(\ell_{it})].$$

In particular,

$$(3.16) \quad \ell_s(\ell_{it}) = f(t, s) \cot \ell_{t+s}$$

$$(3.17) \quad \alpha_t \cdot \alpha_s \cdot \beta_t = f^2(t, s) \cdot \alpha_{t+s}$$

where $f(t, s)$ is holomorphic on t and s . From 3.16 one concludes that $\ell_{0t} \in A_1(x_1, x_3)$, which proves **iii)**. $\square$

Theorem 3.26. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$. Then, up to linear conjugation φ_t is of one of the following types:*

- I)** $\varphi_t[x_0, x_1, x_2, x_3] = [x_0(x_3 + tx_0), e^{\alpha t}x_1x_3 + q_t(x_0, x_2, x_3), e^{\beta t}x_2(x_3 + tx_0), (x_3 + tx_0)^2]$ where:
a) $\alpha = \beta = 0$,

$$q_t(x_0, x_2, x_3) = \frac{c_{33}}{3}t^3x_0^2 + c_{22}tx_2^2 + c_{33}tx_3^2 + \frac{c_{23}}{2}t^2x_0x_2 + c_{33}t^2x_0x_3 + c_{23}tx_2x_3$$

and either $c_{22}, c_{23}, c_{33} \in \{0, 1\}$ or $c_{22} = c_{33} = 1$ and $c_{23} \in \mathbb{C}^*$.

- b)** $\alpha = 0, \beta \neq 0$,

$$q_t(x_0, x_2, x_3) = \frac{c_{33}}{3}t^3x_0^2 + c_{22}(1 - e^{2\beta t})x_2^2 + c_{33}tx_3^2 + (1 - e^{\beta t})c_{02}x_0x_2 + c_{33}t^2x_0x_3$$

and $c_{22}, c_{33}, c_{02} \in \mathbb{C}$.

- c)** $\alpha \neq 0, \beta = 0$,

$$q_t(x_0, x_2, x_3) = (e^{\alpha t} - 1)(c_{00}x_0^2 + c_{22}x_2^2 + c_{02}x_0x_2)$$

and $(c_{00}, c_{22}, c_{02}) = (0, 0, 0), (1, 0, 0), (0, 0, 1)$ or $(c_{00}, 1, 0)$ with $c_{00} \in \mathbb{C}^*$.

- d)** $\alpha \neq 0, \beta \neq 0$,

$$q_t(x_0, x_2, x_3) = c_{00}(e^{\alpha t} - 1)x_0^2 + c_{22}(e^{\alpha t} - e^{2\beta t})x_2^2 + c_{02}(e^{\alpha t} - e^{\beta t})x_0x_2$$

and $c_{00}, c_{22}, c_{02} \in \mathbb{C}$.

- II)** $\varphi_t[x_0, x_1, x_2, x_3] = [x_0(x_3 + tx_0), e^{\alpha t}x_1x_3 + q_t(x_0, x_2, x_3), (x_2 + tx_3 + \frac{t^2}{2}x_0)(x_3 + tx_0), (x_3 + tx_0)^2]$ where
e) $\alpha = 0$,

$$q_t(x_0, x_2, x_3) = t(c_{22}x_2 + c_{02}x_0 + c_{23}x_3)x_2 + \left(\frac{c_{02}}{6}t^3 + \frac{c_{23}}{8}t^4 + \frac{c_{22}}{20}t^5\right)x_0^2 \\ + \left(\frac{c_{23}}{2}t^2 + \frac{c_{22}}{3}t^3\right)(x_3^2 + x_0x_2) + \left(\frac{c_{02}}{2}t^2 + \frac{c_{23}}{2}t^3 + \frac{c_{22}}{4}t^4\right)x_0x_3 + c_{22}t^2x_2x_3$$

and $(c_{02}, c_{22}, c_{23}) = (0, 1, c_{23}), (c_{02}, 0, 0)$ or $(c_{02}, 0, 1)$ with $c_{23}, c_{02} \in \mathbb{C}$.

- f)** $\alpha \neq 0$,

$$q_t(x_0, x_2, x_3) = (e^{\alpha t} - 1)(c_{00}x_0^2 + c_{22}x_2^2 + c_{02}x_0x_2) - \left(\frac{c_{22}}{4}t^4 + \frac{c_{02}}{2}t^2\right)x_0^2 \\ - c_{22}t^2(x_3^2 + x_0x_2) - 2c_{22}tx_2x_3 - (c_{02}t + c_{22}t^3)x_0x_3$$

and $(c_{00}, c_{22}, c_{02}) = (0, 0, 1), (1, 0, 0)$ or $(c_{00}, 0, 1)$ with $c_{00} \in \mathbb{C}$.

Moreover two such different flows φ_t and $\tilde{\varphi}_t$ are linearly conjugated if and only if we are in cases **b)** or **d)** and there exists $\lambda \in \mathbb{C}^*$ such that $\tilde{q}_t = \lambda \cdot q_t$,

Proof. We will use the same notation as in theorem 3.24. Recall that we had chosen coordinates such that $P = P_I = [0, 1, 0, 0]$, $L_I = (x_0 = x_3 = 0)$, $S = (x_3 = 0)$ and $H_t = (a(t)x_0 + x_3 = 0)$. Moreover, we had concluded that

$$\varphi_t[x_0, x_1, x_2, x_3] = [\ell_{0t}\ell_t, \alpha_tx_1x_3 + q_t(x_0, x_2, x_3), \ell_{2t}\ell_t, \beta_t\ell_t^2]$$

with $\ell_{0t} \in A_1(x_0, x_3)$, $\ell_{2t} \in A_1(x_0, x_2, x_3)$ such that

$$\alpha_0 = 1, q_0 = 0, \ell_{00} = x_0, \ell_{20} = x_2, \ell_{30} = x_3, \ell_{10} = x_3, a(0) = 0$$

and

$$(3.18) \quad \ell_s(\ell_{it}) = f(t, s) \cdot \ell_{t+s},$$

$$(3.19) \quad \alpha_t \cdot \alpha_s \cdot \beta_t = f^2(t, s) \cdot \alpha_{t+s}.$$

Moreover,

$$\varphi_s(\varphi_t(x)) = [\ell_{0s}(\ell_{it})\ell_s(\ell_{it}), \alpha_s\alpha_t\beta_tx_1x_3 + \beta_t\alpha_sq_t(x_0, x_2, x_3) + q_s(\ell_{it}), \ell_{2s}(\ell_{it})\ell_s(\ell_{it}), \beta_t\ell_s^2(\ell_{it})]$$

and ℓ_t is a common factor of all the components of $\varphi_s \circ \varphi_t$.

We denote now $\ell_{0t} = A_0^0(t)x_0 + A_0^3(t)x_3$. We can rewrite 3.18 as

$$(a(s)A_0^0(t) + a(t))x_0 + (a(s)A_0^3(t) + 1)x_3 = f(t, s)(a(t+s)x_0 + x_3)$$

so

$$\begin{aligned} a(s)A_0^0(t) + a(t) &= f(t, s)a(t+s) \\ a(s)A_0^3(t) + 1 &= f(t, s). \end{aligned}$$

As $a(0) = 0$ we have $f(t, 0) = 1$. On the other hand, as $\alpha_0 = 1$ one has

$$\alpha_t\beta_t = f^2(t, 0)\alpha_t$$

so $\beta_t = 1$. We can define $\psi_t[x_0, x_2, x_3] = [\ell_{0t}, \ell_{2t}, \ell_t]$, which is a linear flow. Moreover, $\ell_t = a(t)x_0 + x_3$ and $\ell_{0t} \in A_1(x_0, x_3)$. From the classification of linear flows on $\mathbb{P}^2$ we derive that either

$$a(t) = t, \ell_{0s} = x_0, \ell_{2s} = x_2e^{\beta t}$$

or

$$a(t) = t, \ell_{0s} = x_0, \ell_{2s} = x_2 + tx_3 + \frac{t^2}{2}x_0,$$

which in both cases yields $f(t, s) = 1$ and $\alpha_t = e^{\alpha t}$ with $\alpha, \beta \in \mathbb{C}$. Thus, either

$$\varphi_t[x_0, x_1, x_2, x_3] = [x_0(x_3 + tx_0), e^{\alpha t}x_1x_3 + q_t(x_0, x_2, x_3), e^{\beta t}x_2(x_3 + tx_0), (x_3 + tx_0)^2]$$

and

$$(3.20) \quad \varphi_s(\varphi_t)(x) = [x_0(x_3 + (t+s)x_0), e^{\alpha(t+s)}x_1x_3 + e^{\alpha s}q_t(x) + q_t(\ell_{it}), e^{\beta(t+s)}x_2(x_3 + (t+s)x_0), (x_3 + (t+s)x_0)^2]$$

or

$$\varphi_t[x_0, x_1, x_2, x_3] = \left[x_0(x_3 + tx_0), e^{\alpha t}x_1x_3 + q_t(x_0, x_2, x_3), \left(x_2 + tx_3 + \frac{t^2}{2}x_0 \right) (x_3 + tx_0), (x_3 + tx_0)^2 \right]$$

and

$$(3.21) \quad \varphi_s(\varphi_t)(x) = \left[x_0(x_3 + (t+s)x_0), e^{\alpha(t+s)}x_1x_3 + e^{\alpha s}q_t(x) + q_t(\ell_{it}), \left(x_2 + (t+s)x_3 + \frac{(t+s)^2}{2}x_0 \right) (x_3 + (t+s)x_0), (x_3 + (t+s)x_0)^2 \right].$$

We must impose now that either

$$(3.22) \quad e^{\alpha s}q_t(x_0, x_2, x_3) + q_s(x_0, e^{\beta t}x_2, x_3 + tx_0) = q_{t+s}(x_0, x_2, x_3)$$

or

$$(3.23) \quad e^{\alpha s}q_t(x_0, x_2, x_3) + q_s(x_0, x_2 + tx_3 + \frac{t^2}{2}x_0, x_3 + tx_0) = q_{t+s}(x_0, x_2, x_3).$$

We will denote

$$q_t(x_0, x_2, x_3) = q_{00}(t)x_0^2 + q_{02}(t)x_0x_2 + q_{03}(t)x_0x_3 + q_{22}(t)x_2^2 + q_{23}(t)x_2x_3 + q_{33}(t)x_3^2.$$

From 3.22 we obtain

$$\begin{aligned}
q_{00}(t+s) &= e^{\alpha s} q_{00}(t) + q_{00}(s) + t^2 q_{33}(s) + t q_{03}(s), \\
q_{02}(t+s) &= e^{\alpha s} q_{02}(t) + e^{\beta t} (q_{02}(s) + t q_{23}(s)), \\
q_{03}(t+s) &= e^{\alpha s} q_{03}(t) + q_{03}(s) + 2t q_{33}(s), \\
q_{22}(t+s) &= e^{\alpha s} q_{22}(t) + e^{2\beta t} q_{22}(s), \\
q_{23}(t+s) &= e^{\alpha s} q_{23}(t) + e^{\beta t} q_{23}(s), \\
q_{33}(t+s) &= e^{\alpha s} q_{33}(t) + q_{33}(s).
\end{aligned}$$

Using the symmetry on t, s on the previous equations one concludes that $q_t[x_0, x_2, x_3]$ verifies 3.22 if and only if it is of one forms:

- a) $(c_{00}t + \frac{c_{03}}{2}t^2 + \frac{c_{33}}{3}t^3)x_0^2 + c_{22}tx_2^2 + c_{33}tx_3^2 + (c_{02}t + \frac{c_{23}}{2}t^2)x_0x_2 + (c_{03}t + c_{33}t^2)x_0x_3 + c_{23}tx_2x_3$, for $\alpha = \beta = 0$,
- b) $(c_{00}t + \frac{c_{03}}{2}t^2 + \frac{c_{33}}{3}t^3)x_0^2 + c_{22}(1 - e^{2\beta t})x_2^2 + c_{33}tx_3^2 + ((1 - e^{\beta t})c_{02} - c_{23}te^{\beta t})x_0x_2 + (c_{03}t + c_{33}t^2)x_0x_3 + c_{23}(1 - e^{\beta t})x_2x_3$, for $\alpha = 0, \beta \neq 0$, or
- c) $(c_{00}(e^{\alpha t} - 1) - c_{03}t - c_{33}t^2)x_0^2 + c_{22}(e^{\alpha t} - 1)x_2^2 + c_{33}(e^{\alpha t} - 1)x_3^2 + (c_{02}(e^{\alpha t} - 1) - c_{23}t)x_0x_2 + (c_{03}(e^{\alpha t} - 1) - 2c_{33}t)x_0x_3 + c_{23}(e^{\alpha t} - 1)x_2x_3$ for $\alpha \neq 0$ and $\beta = 0$,
- d) $(c_{00}(e^{\alpha t} - 1) - c_{03}t - c_{33}t^2)x_0^2 + c_{22}(e^{\alpha t} - e^{2\beta t})x_2^2 + c_{33}(e^{\alpha t} - 1)x_3^2 + (c_{02}(e^{\alpha t} - e^{\beta t}) - c_{23}te^{\beta t})x_0x_2 + (c_{03}(e^{\alpha t} - 1) - 2c_{33}t)x_0x_3 + c_{23}(e^{\alpha t} - e^{\beta t})x_2x_3$ for $\alpha \neq 0, \beta \neq 0$,

and c_{ij} are constant. Note for instance that the last equation yields

$$e^{\alpha s} q_{33}(t) + q_{33}(s) = e^{\alpha t} q_{33}(s) + q_{33}(t)$$

which implies that $\frac{q_{33}(t)}{(e^{\alpha t} - 1)}$ is constant for $\alpha \neq 0$.

A similar computation allows to conclude that q_t verifies equation 3.23 if and only if $q_t(x_0, x_2, x_3)$ is

- e) $tq_0 + (\frac{c_{03}}{2}t^2 + (c_{33} + \frac{c_{02}}{2})\frac{t^3}{3} + \frac{c_{23}}{8}t^4 + \frac{c_{22}}{20}t^5)x_0^2 + (\frac{c_{23}}{2}t^2 + \frac{c_{22}}{3}t^3)x_3^2 + (\frac{c_{23}}{2}t^2 + \frac{c_{22}}{3}t^3)x_0x_2 + ((c_{33} + \frac{c_{02}}{2})t^2 + \frac{c_{23}}{2}t^3 + \frac{c_{22}}{4}t^4)x_0x_3 + c_{22}t^2x_2x_3$, for $\alpha = 0$
- f) $(e^{\alpha t} - 1)q_0 - (\frac{c_{22}}{4}t^4 + \frac{c_{23}}{2}t^3 + (c_{33} + \frac{c_{02}}{2})t^2 + c_{03}t)x_0^2 - (tc_{23} + t^2c_{22})(x_3^2 + x_0x_2) - 2tc_{22}x_2x_3 - (tc_{02} + 2tc_{33} + \frac{3c_{23}}{2}t^2 + c_{22}t^3)x_0x_3$, for $\alpha \neq 0$.

where

$$q_0(x_0, x_2, x_3) = c_{00}x_0^2 + c_{22}x_2^2 + c_{33}x_3^2 + c_{02}x_0x_2 + c_{03}x_0x_3 + c_{23}x_2x_3$$

and c_{ij} are constant.

Let now φ_t and $\tilde{\varphi}_t$ be two flows of one of the previous types: **a)**, ..., **f)**, and $A = [\mu_0, \mu_1, \mu_2, \mu_3]$ with $\mu_i \in A_1(x_0, x_1, x_2, x_3)$ for $i = 0, 1, 2, 3$ a linear conjugation such that

$$A \circ \varphi_t = \tilde{\varphi}_t \circ A. \quad (*)$$

Using that $P_I = [0, 1, 0, 0]$, $L_I = (x_0 = x_3 = 0)$ and $S = (x_3^2 = 0)$ are fix and common for both flows, $H_t = (tx_0 + x_3 = 0)$ and imposing $(*)$ on the last component we conclude that

$$\mu_0 = x_0; \quad \mu_1 = \mu_1^0 x_0 + \mu_1^1 x_1 + \mu_1^2 x_2 + \mu_1^3 x_3; \quad \mu_2 = \mu_2^0 x_0 + \mu_2^2 x_2 + \mu_2^3 x_3; \quad \mu_3 = x_3,$$

up to product by an element of $\mathbb{C}^*$.

If φ_t and $\tilde{\varphi}_t$ are both of type **I)** then imposing now $(*)$ on the third component of the flows one obtains

$$\mu_2^0 e^{\tilde{\beta} t} x_0 + \mu_2^2 e^{\tilde{\beta} t} x_2 + \mu_2^3 e^{\tilde{\beta} t} x_3 = \mu_2^0 x_0 + \mu_2^2 e^{\beta t} x_2 + \mu_2^3 (x_3 + tx_0).$$

As $\mu_2^2 \neq 0$ we conclude that $\beta = \tilde{\beta}$ and $\mu_2^3 = 0$. Moreover either β or μ_2^0 are zero and if $\mu_2^0 = 0$ then $\mu_2^2 = 1$.

If both flows are of type **II)** analogously one obtains $\mu_2^3 = 0$ and $\mu_2^2 = 1$.

Finally, it also implies that one flow of type **I)** and one flow of type **II)** cannot be conjugated.

Imposing condition $(*)$ on the second component for two linearly conjugated flows of either type **I)** or two flows of type **II)** one concludes easily that in both cases $\alpha = \tilde{\alpha}$. Moreover, writing explicitly this condition for the six cases **a)** to **f)** one obtains the list of the statement. Let us assume for instance that $\alpha = \beta = 0$, one obtains:

$$\begin{aligned}
 & (\mu_1^0 x_0 + \mu_1^1 x_1 + \mu_2^0 x_2 + \mu_1^3 x_3) x_3 + (\tilde{c}_{00} t + \frac{\tilde{c}_{03}}{2} t^2 + \frac{\tilde{c}_{33}}{3} t^3) x_0^2 + (\tilde{c}_{02} t + \frac{\tilde{c}_{23}}{2} t^2) x_0 (\mu_2^0 x_0 + \mu_2^2 x_2) + \\
 & \tilde{c}_{22} t (\mu_2^0 x_0 + \mu_2^2 x_2)^2 + \tilde{c}_{23} t (\mu_2^0 x_0 + \mu_2^2 x_2) x_3 + \tilde{c}_{33} t x_3^2 + (\tilde{c}_{03} t + \tilde{c}_{33} t^2) x_0 x_3 = \\
 & \mu_1^0 x_0 (x_3 + t x_0) + \mu_1^1 (x_1 x_3 + (c_{00} t + \frac{c_{03}}{2} t^2 + \frac{c_{33}}{3} t^3) x_0^2 + (c_{02} t + \frac{c_{23}}{2} t^2) x_0 x_2 + c_{22} t x_2^2 + c_{23} t x_2 x_3 + c_{33} t x_3^2 + \\
 & (c_{03} t + c_{33} t^2) x_0 x_3) + \mu_1^2 x_2 (x_3 + t x_0) + \mu_1^3 (x_3 + t x_0)^2,
 \end{aligned}$$

which implies

$$\begin{aligned}
 \tilde{c}_{00} &= \mu_1^1 c_{00} - \mu_2^0 \mu_1^1 (\mu_2^2)^{-1} c_{02} + \mu_2^0 (\mu_2^2)^{-2} \mu_1^1 c_{22} - \mu_2^0 \mu_1^2 (\mu_2^2)^{-1} + \mu_1^0 \\
 \tilde{c}_{22} &= (\mu_2^2)^{-2} \mu_1^1 c_{22} \\
 \tilde{c}_{33} &= \mu_1^1 c_{33} \\
 \tilde{c}_{02} &= (\mu_2^2)^{-1} \mu_1^1 c_{02} - 2 \mu_2^0 (\mu_2^2)^{-2} \mu_1^1 c_{22} + \mu_1^2 (\mu_2^2)^{-1} \\
 \tilde{c}_{03} &= \mu_1^1 c_{03} - \mu_2^0 (\mu_2^2)^{-1} \mu_1^1 c_{23} + 2 \mu_1^3 \\
 \tilde{c}_{23} &= (\mu_2^2)^{-1} \mu_1^1 c_{23}.
 \end{aligned}$$

It is not difficult to see that up to a linear conjugation one can choose the parameters c_{ij} as in the statement. And analogous computation gives the rest of the flows. $\square$

Remark 3.27. Computing $\text{Ind}(\varphi_t)$ it is not difficult to verify that the flows in 3.26 belong indeed to $\mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{tan}^{[2]}(\times) \cup \mathbf{lin}$

Remark 3.28. As in the generic case, up to a normalisation of the time coordinate t by an homotecy one can assume that α is either 0 or 1.

3.3. Flows in $\mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$.

Definition 3.29. Let φ_t be a quadratic flow. We say that φ_t is a *polynomial flow* if there is an invariant chart $\mathbb{C}^3$ such that $\varphi_t|_{\mathbb{C}^3} : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ is polynomial for each t .

The following result was proved by D. Cerveau and J. Deserti for quadratic flows in $\mathbb{P}^2$. The proof for $\mathbb{P}^3$ is almost identical, we include it for the sake of clarity.

Proposition 3.30. *Let φ_t be a quadratic flow such that H_t is fix. Then φ_t is a polynomial flow.*

Proof. We can assume that φ_t is not linear and that $H = (x_3 = 0)$. Therefore for every t such that φ_t is quadratic we can write the flow as

$$A_t(x_0 x_3, x_1 x_3 - x_2^2, x_2 x_3, x_3^2) B_t$$

or

$$A_t(x_0 x_3 - x_1 x_2, x_1 x_2, x_2 x_3, x_3^2) B_t$$

where $A_t, B_t \in \text{PGL}_4(\mathbb{C})$ and B_t preserves the hyperplan $x_3 = 0$. As $\text{Ind}(\varphi_t)$ is contained in H , i.e. at the infinity of the affine chart $x_3 = 1$ for almost every t and thus for every t , we have that φ_t is polynomial. $\square$

Lemma 3.31. *Let φ_t a quadratic flow in $\mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$. Then P_I is fix.*

Proof. Assume that $\varphi_t \in \mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$ and that P_{I_t} is mobile. We will show that $\varphi_t \in \mathbf{tan}^{[2]}(\mathbf{O}) \cup \mathbf{lin}$. By lemma 3.8 we have that H_t is fix. Without loss of generality we can assume that $H = (x_3 = 0)$ and $P_t = P_{I-t} = [\alpha_t, \beta_t, \gamma_t, 0]$ (because $P_{I_t} \in H$). Therefore

$$\varphi_t(x) = [\alpha_t q_t + \ell_{0t} x_3, \beta_t q_t + \ell_{1t} x_3, \gamma_t q_t + \ell_{2t} x_3, \ell_{3t} x_3]$$

with $\ell_{it} \in A_1(x_0, x_1, x_2, x_3)$ for $i = 0, 1, 2, 3$, $q_t \in A_2(x_0, x_1, x_2)$, $\text{rank } q_t \leq 2$ and $\text{Jac}(\varphi_t) = f(t) x_3^4$ with $f(t) \in \mathbb{C}^*$. Moreover, by the previous lemma φ_t is a polynomial flow. In the affine coordinates $x = \frac{x_0}{x_3}$, $y = \frac{x_1}{x_3}$ and $z = \frac{x_2}{x_3}$, the flow φ_t has the expression:

$$\tilde{\varphi}_t(x) = \left(\frac{\alpha_t \frac{q_t}{x_3} + \frac{\ell_{0t}}{x_3}}{\frac{\ell_{3t}}{x_3}}, \frac{\beta_t \frac{q_t}{x_3} + \frac{\ell_{1t}}{x_3}}{\frac{\ell_{3t}}{x_3}}, \frac{\gamma_t \frac{q_t}{x_3} + \frac{\ell_{2t}}{x_3}}{\frac{\ell_{3t}}{x_3}} \right).$$

As $\tilde{\varphi}_t$ is a polynomial we must have $\ell_{3t} = \delta_t x_3$ so (dividing by δ_t if necessary) we have $\ell_{3t} = x_3$. Thus, in affine coordinates

$$\tilde{\varphi}_t = (\alpha_t \tilde{q}_t + \tilde{\ell}_{0t}, \beta_t \tilde{q}_t + \tilde{\ell}_{1t}, \gamma_t \tilde{q}_t + \tilde{\ell}_{2t}),$$

where $\tilde{\ell}_{it} = A_t^i x + B_t^i y + C_t^i z + D_t^i$ with $A_t^i, B_t^i, C_t^i, D_t^i \in \mathbb{C}$ for $i = 0, 1, 2$ and $\tilde{q}_t$ is a homogeneous polynomial of degree 2 in x, y, z . On the other hand $\tilde{\varphi}_s \circ \tilde{\varphi}_t$ must still be a quadratic flow. Let $A_{\tilde{q}_t}$ be the bilinear form associated to $\tilde{q}_t$, then denoting $\tilde{L}_t = (\tilde{\ell}_{0t}, \tilde{\ell}_{1t}, \tilde{\ell}_{2t})$ one has

$$\tilde{q}_s(\tilde{\varphi}_t(x, y, z)) = (\tilde{q}_t P_t + \tilde{L}_t) \cdot A_{\tilde{q}_s} \cdot (\tilde{q}_t P_t + \tilde{L}_t)^t = \tilde{q}_t^2(P_t \cdot A_{\tilde{q}_s} \cdot P_t^T) + 2\tilde{q}_t(P_t \cdot A_{\tilde{q}_s} \cdot \tilde{L}_t^T) + \tilde{L}_t \cdot A_{\tilde{q}_s} \cdot \tilde{L}_t^T.$$

As the two first terms have degrees 4 and 3 respectively they must be zero. We conclude that

$$(3.24) \quad P_t \cdot A_{\tilde{q}_s} \cdot P_t^T = \tilde{q}_s(P_t) = 0$$

$$(3.25) \quad P_t \cdot A_{\tilde{q}_s} \cdot \tilde{L}_t^T = 0$$

As P_t is mobile condition 3.24 implies that $\tilde{q}_s$ has a fixed component. We can assume that $q_t = x_1 \cdot m_t$ where $m_t \in A_1(x_0, x_1, x_2)$ and $\beta_t = 0$, i.e. $P_t = (\alpha_t, 0, \gamma_t)$. Thus

$$\tilde{\varphi}_t(x, y, z) = (\alpha_t \cdot y \cdot \tilde{m}_t + \tilde{\ell}_{0t}, \gamma_t \cdot y \cdot \tilde{m}_t + \tilde{\ell}_{2t}),$$

where $\tilde{m}_t$ is the homogeneous polynomial of degree 1 on x, y, z obtained from m_t . Now, condition 3.25 says that either $\tilde{\ell}_{1t} = 0$ or $\frac{1}{2}\tilde{m}_s(P_t) = P_t \cdot A_{\tilde{q}_s} = 0$. In the first case $\tilde{\varphi}_t$ wouldn't be an automorphism, which contradicts the hypothesis. In the second case, as P_t is mobile, $\tilde{m}_s$ must be fix and in fact a multiple of y . Thus, one can assume

$$\tilde{\varphi}_t(x, y, z) = (\alpha_t y^2 + \tilde{\ell}_{0t}, \gamma_t y^2 + \tilde{\ell}_{2t})$$

and $\varphi_t \in \mathbf{tan}^{[2]}(//) \cup \mathbf{lin}$, which ends the proof. $\square$

Lemma 3.32. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(//) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$. Then H_t is fix.*

Proof. As P_{I_t} mobile implies that H is fix it is enough to consider the case P_I fix. Assume that $H_t = (\ell_t = 0)$ is mobile and $P_I = [1, 0, 0, 0]$. Therefore

$$\varphi_t(x) = [\ell_{0t}\ell_t + q_t, \ell_{1t}\ell_t, \ell_{2t}\ell_t, \ell_{3t}\ell_t],$$

with $\ell_{0t}, \ell_{1t}, \ell_{2t}, \ell_{3t} \in A_1(x_0, x_1, x_2, x_3)$. Note also that $P_I \in (\ell_t = 0)$ and that the strict transform of a line through P_I is a conic through P_I , so in fact $\ell_{0t}, \ell_{1t}, \ell_{2t}, \ell_{3t} \in A_1(x_1, x_2, x_3)$. Moreover $P_I \in \text{Ind}(\varphi_t)$ implies that q_t has no term in x_0^2 . Note that for φ_t to be birational we must have $\ell_{1t}, \ell_{2t}, \ell_{3t}$ linearly independent and note also that we can assume that $\ell_{0t} = \delta_t x_0$ (modifying q_t if necessary). Then

$$\varphi_t[x_0, x_1, x_2, x_3] = [\delta_t x_0 + \frac{q_t(x_1, x_2, x_3)}{\ell_t(x_1, x_2, x_3)}, \ell_{1t}, \ell_{2t}, \ell_{3t}].$$

As $\varphi_0 = Id$ we have $q_0 = 0$, $\delta_0 = 1$ and $\ell_{i0} = x_i$ for $i = 1, 2, 3$. We impose now the condition $\varphi_s \circ \varphi_t = \varphi_{t+s}$. We have

$$\varphi_s \circ \varphi_t(x) = \left[\delta_s \delta_t x_0 + \delta_s \frac{q_t(x)}{\ell_t(x)} + \frac{q_s(\ell_{it})}{\ell_s(\ell_{it})}, \ell_{1s}(\ell_{it}), \ell_{2s}(\ell_{it}), \ell_{3s}(\ell_{it}) \right].$$

In particular $\Psi_t[x_1, x_2, x_3] = [\ell_{1t}, \ell_{2t}, \ell_{3t}]$ is a linear flow in $\mathbb{P}^2$. Thus we can assume that $\delta_t \cdot \delta_s = \delta_{t+s}$ and

$$\delta_s \frac{q_t(x_1, x_2, x_3)}{\ell_t(x_1, x_2, x_3)} + \frac{q_s(\ell_{it})}{\ell_s(\ell_{it})} = \frac{q_{t+s}(x)}{\ell_{t+s}(x)}$$

so

$$(3.26) \quad (\delta_s \cdot q_t \cdot \ell_s(\ell_{it}) + q_s(\ell_{it}) \cdot \ell_t) \cdot \ell_{t+s} = q_{t+s} \cdot \ell_s(\ell_{it}) \cdot \ell_t.$$

As H_t is mobile ℓ_t cannot divide ℓ_{t+s} so it divides the first factor of the left part of equation 3.26, which implies that ℓ_t divides the product $q_t \cdot \ell_s(\ell_{it})$. If ℓ_t divides q_t for all t the flow is linear, which is a contradiction so we conclude that ℓ_t is a non-zero scalar multiple of $\ell_s(\ell_{it})$. A similar argument shows that ℓ_{t+s} is a non-zero scalar multiple of $\ell_s(\ell_{it})$. Again this would imply that H_t is fix. $\square$

Theorem 3.33. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(//) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$. Then:*

- i) H is fix.
- ii) φ_t is a polynomial flow.
- iii) If $\varphi_t \in \mathbf{tan}^{[2]}(//) \cup \mathbf{lin}$ then C_I is fix.

- iv) If $\varphi_t \in \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$ then P_I is fix.
- v) If P_I is fix then φ_t preserves the family of hyperplanes through P_I .
- vi) If C_I is fix then φ_t preserves the family of hyperplanes through a line L contained in C_I .

Moreover, up to linear conjugation, φ_t belongs to the following list:

- I) If P_{It} is mobile and $\varphi_t \in \mathbf{tan}^{[2]}(//) \cup \mathbf{lin}$.
 - a) $\varphi_t(x) = [a(e^{2\mu t} - e^{\alpha t})x_1^2 + e^{\alpha t}x_0x_3, e^{\mu t}x_1x_3, b(e^{2\mu t} - e^{\beta t})x_1^2 + e^{\beta t}x_2x_3, x_3^2]$,
 - b) $\varphi_t(x) = [a(e^{2\mu t} - e^{\mu t})x_1^2 + e^{\mu t}(x_0 + tx_1)x_3, e^{\mu t}x_1x_3, b(e^{2\mu t} - e^{\beta t})x_1^2 + e^{\beta t}x_2x_3, x_3^2]$,
 - c) $\varphi_t(x) = [a(e^{2\mu t} - e^{\alpha t})x_1^2 + e^{\alpha t}x_0x_3, e^{\mu t}x_1x_3, b(e^{2\mu t} - 1)x_1^2 + x_2x_3 + tx_3^2, x_3^2]$,
 - d) $\varphi_t(x) = [a(e^{2\mu t} - e^{\mu t})x_1^2 + e^{\mu t}(x_0 + tx_1)x_3, e^{\mu t}x_1x_3, b(e^{2\mu t} - 1)x_1^2 + x_2x_3 + tx_3^2, x_3^2]$.

with $a, b \neq 0$.
- II) If P_I is fix then:
 - 1) $\varphi_t(x) = [e^{\delta t}x_0x_3 + 2tBe^{\alpha t}x_1x_3 + 2tCx_2x_3 + t^2Cx_3^2 + A(e^{2\alpha t} - e^{\delta t})x_1^2 + 2B(e^{\alpha t} - e^{\delta t})x_1x_2 + C(1 - e^{\delta t})x_2^2, e^{\alpha t}x_1x_3, x_2x_3 + tx_3^2, x_3^2]$ with $\delta \neq 0, \delta \neq \alpha, \delta \neq 2\alpha$,
 - 2) $\varphi_t(x) = [x_0x_3 + 2tBe^{\alpha t}x_1x_3 + (bt + Ct^2)x_2x_3 + (\frac{b}{2}t^2 + \frac{C}{3}t^3)x_3^2 + A(e^{2\alpha t} - 1)x_1^2 + 2B(e^{\alpha t} - 1)x_1x_2, e^{\alpha t}x_1x_3, x_2x_3 + tx_3^2, x_3^2]$ with $\alpha \neq 0$,
 - 3) $\varphi_t(x) = [x_0x_3 + (at + Bt^2)x_1x_3 + (bt + Ct^2)x_2x_3 + (\frac{b}{2}t^2 + \frac{C}{3}t^3)x_3^2 + Atx_1^2 + 2Btx_1x_2 + Ctx_2^2, x_1x_3, x_2x_3 + tx_3^2, x_3^2]$,
 - 4) $\varphi_t(x) = [e^{\delta t}x_0x_3 + (at + Bt^2)e^{\delta t}x_1x_3 + 2Ctx_2x_3 + Ct^2x_3^2 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2Bte^{\delta t}x_1x_2 + C(1 - e^{\delta t})x_2^2, e^{\delta t}x_1x_3, x_2x_3 + tx_3^2, x_3^2]$ with $\delta \neq 0$,
 - 5) $\varphi_t(x) = [e^{\delta t}x_0x_3 + 2tBe^{\frac{\delta t}{2}}x_1x_3 + 2Ctx_2x_3 + Ct^2x_3^2 + Ate^{\delta t}x_1^2 + 2B(e^{\frac{\delta t}{2}} - e^{\delta t})x_1x_2 + C(1 - e^{\delta t})x_2^2, e^{\frac{\delta t}{2}}x_1x_3, x_2x_3 + tx_3^2, x_3^2]$ with $\delta \neq 0$,
 - 6) $\varphi_t(x) = [e^{\delta t}x_0x_1 + A(e^{-\alpha t} - e^{\delta t})x_2^2 + 2(B(e^{-\alpha t} - e^{\delta t}) + Ate^{-\alpha t})x_2x_3 + (C(e^{-\alpha t} - e^{\delta t}) + (At^2 + 2tB)e^{-\alpha t})x_3^2, e^{\alpha t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\delta \neq 0$,
 - 7) $\varphi_t(x) = [e^{\delta t}x_0x_1 + Ate^{\delta t}x_2^2 + (2Bt + At^2)e^{\delta t}x_2x_3 + (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t}x_3^2, e^{-\delta t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\delta \neq 0$,
 - 8) $\varphi_t(x) = [e^{\delta t}x_0x_1 + ate^{\delta t}x_1^2 + A(e^{-\delta t} - e^{\delta t})x_2^2 + 2(B(e^{-\delta t} - e^{\delta t}) + Ate^{-\delta t})x_2x_3 + (C(e^{-\delta t} - e^{\delta t}) + (At^2 + 2tB)e^{-\delta t})x_3^2, e^{\delta t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\delta \neq 0$,
 - 9) $\varphi_t(x) = [x_0x_1 + btx_1x_2 + \frac{bt^2}{2}x_1x_3 + A(e^{-\alpha t} - 1)x_2^2 + 2(B(e^{-\alpha t} - 1) + Ate^{-\alpha t})x_2x_3 + (C(e^{-\alpha t} - 1) + (At^2 + 2tB)e^{-\alpha t})x_3^2, e^{\alpha t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\alpha \neq 0$,
 - 10) $\varphi_t(x) = [x_0x_1 + atx_1^2 + btx_1x_2 + \frac{bt^2}{2}x_1x_3 + Atx_2^2 + 2(Bt + \frac{A}{2}t^2)x_2x_3 + (Ct + Bt^2 + \frac{A}{3}t^3)x_3^2, x_1^2, (x_2 + tx_3)x_1, x_3x_1]$,
 - 11) $\varphi_t(x) = [e^{\delta t}x_0x_3 + A(e^{2\alpha t} - e^{\delta t})x_1^2 + 2B(e^{(\alpha+\beta)t} - e^{\delta t})x_1x_2 + C(e^{2\beta t} - e^{\delta t})x_2^2, e^{\alpha t}x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\delta \neq 0, \delta \neq \alpha, \beta, \alpha + \beta, 2\alpha, 2\beta$,
 - 12) $\varphi_t(x) = [x_0x_3 + ct x_3^2 + A(e^{2\alpha t} - 1)x_1^2 + 2B(e^{(\alpha+\beta)t} - 1)x_1x_2 + C(e^{2\beta t} - 1)x_2^2, e^{\alpha t}x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\alpha \neq 0$ and $\beta \neq 0$,
 - 13) $\varphi_t(x) = [e^{\delta t}x_0x_3 + ate^{\delta t}x_1x_3 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2B(e^{(\delta+\beta)t} - e^{\delta t})x_1x_2 + C(e^{2\beta t} - e^{\delta t})x_2^2, e^{\delta t}x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\delta \neq 0, \delta \neq \beta, 2\beta$,
 - 14) $\varphi_t(x) = [e^{2\alpha t}x_0x_3 + Ate^{2\alpha t}x_1^2 + 2B(e^{(\alpha+\beta)t} - e^{2\alpha t})x_1x_2 + C(e^{2\beta t} - e^{2\alpha t})x_2^2, e^{\alpha t}x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\alpha \neq 0, \beta \neq 0, \alpha \neq \beta$,
 - 15) $\varphi_t(x) = [e^{\delta t}x_0x_3 + ate^{\delta t}x_1x_3 + bte^{\delta t}x_2x_3 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2B(e^{2\delta t} - e^{\delta t})x_1x_2 + C(e^{2\delta t} - e^{\delta t})x_2^2, e^{\delta t}x_1x_3, e^{\delta t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
 - 16) $\varphi_t(x) = [e^{2\alpha t}(x_0x_3 + Atx_1^2 + 2Btx_1x_2 + Ctx_2^2), e^{\alpha t}x_1x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,
 - 17) $\varphi_t(x) = [e^{(\alpha+\beta)t}x_0x_3 + A(e^{2\alpha t} - e^{(\alpha+\beta)t})x_1^2 + 2Bte^{(\alpha+\beta)t}x_1x_2 + C(e^{2\beta t} - e^{(\alpha+\beta)t})x_2^2, e^{\alpha t}x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\alpha \neq 0, \beta \neq 0, \alpha \neq \beta$,
 - 18) $\varphi_t(x) = [e^{\delta t}x_0x_3 + ate^{\delta t}x_1x_3 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2Bte^{\delta t}x_1x_2 + C(1 - e^{\delta t})x_2^2, e^{\delta t}x_1x_3, x_2x_3, x_3^2]$ with $\delta \neq 0$,
 - 19) $\varphi_t(x) = [x_0x_3 + atx_1x_3 + ct x_3^2 + Atx_1^2 + 2B(e^{\beta t} - 1)x_1x_2 + C(e^{2\beta t} - 1)x_2^2, x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\beta \neq 0$,
 - 20) $\varphi_t(x) = [x_0x_3 + t(cx_3^2 + \epsilon_1x_1^2 + \epsilon_2x_2^2), x_1x_3, x_2x_3, x_3^2]$, with $\epsilon_i = 0$ or 1 for $i = 1, 2$,
 - 21) $\varphi_t(x) = [e^{\delta t}x_0x_3 + 2Atx_1x_3 + (2Bte^{\alpha t} - 2At)x_2x_3 + At^2x_3^2 + A(1 - e^{\delta t})x_1^2 + 2(B(e^{\alpha t} - e^{\delta t}) - A(1 - e^{\delta t}))x_1x_2 + (C(e^{2\alpha t} - e^{\delta t}) - 2B(e^{\alpha t} - e^{\delta t}) + A(1 - e^{\delta t}))x_2^2, x_1x_3 + (e^{\alpha t} - 1)x_2x_3 + tx_3^2, e^{\alpha t}x_2x_3, x_3^2]$ with $\delta \neq 0, \alpha, 2\alpha, \alpha \neq 0$,

- 22) $\varphi_t(x) = [e^{\delta t}x_0x_3 + 2Atx_1x_3 + ((Bt^2 + bt)e^{\delta t} - 2At)x_2x_3 + At^2x_3^2 + A(1 - e^{\delta t})x_1^2 + 2(Bte^{\delta t} - A(1 - e^{\delta t}))x_1x_2 + (C(e^{2\delta t} - e^{\delta t}) - 2Bte^{\delta t} + A(1 - e^{\delta t}))x_2^2, x_1x_3 + (e^{\delta t} - 1)x_2x_3 + tx_3^2, e^{\delta t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
- 23) $\varphi_t(x) = [e^{2\alpha}x_0x_3 + 2tAx_1x_3 + (2te^{\alpha t}B - 2tA)x_2x_3 + At^2x_3^2 + A(1 - e^{2\alpha t})x_1^2 + 2(B(e^{\alpha t} - e^{2\alpha t}) - A(1 - e^{2\alpha t}))x_1x_2 + (Cte^{2\alpha t} - 2B(e^{\alpha t} - e^{2\alpha t}) + A(1 - e^{2\alpha t}))x_2^2, x_1x_3 + (e^{\alpha t} - 1)x_2x_3 + tx_3^2, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,
- 24) $\varphi_t(x) = [x_0x_3 + (at + At^2)x_1x_3 + (2te^{\alpha t}B - At^2 - at)x_2x_3 + (\frac{A}{3}t^3 + \frac{a}{2}t^2)x_3^2 + Atx_1^2 + 2(B(e^{\alpha t} - 1) - At)x_1x_2 + (C(e^{2\alpha t} - 1) - 2B(e^{\alpha t} - 1) + At)x_2^2, x_1x_3 + (e^{\alpha t} - 1)x_2x_3 + tx_3^2, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,
- 25) $\varphi_t(x) = [e^{\delta t}x_0x_2 - 2A(e^{-\alpha t} - e^{\delta t})x_1x_2 + A(e^{-\alpha t} - e^{\delta t})x_2^2 + (-2Ate^{-\alpha t} + 2B(1 - e^{-\alpha t}))x_2x_3 + A(e^{-\alpha t} - e^{\delta t})x_1^2 + 2(B(e^{-\alpha t} - e^{\delta t}) + Ate^{-\alpha t})x_1x_3 + (C(e^{-\alpha t} - e^{\delta t}) + At^2e^{-\alpha t} + 2Bte^{-\alpha t})x_3^2, (x_1 + (e^{\alpha t} - 1)x_2 + tx_3)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\delta \neq 0, \alpha \neq 0$,
- 26) $\varphi_t(x) = [x_0x_2 + (at - 2A(e^{-\alpha t} - 1))x_1x_2 + (-at + A(e^{-\alpha t} - 1))x_2^2 + (\frac{a}{2}t^2 - 2Ate^{-\alpha t} + 2(1 - e^{-\alpha t})B)x_2x_3 + A(e^{-\alpha t} - e^{\delta t})x_1^2 + 2(B(e^{-\alpha t} - 1) + Ate^{-\alpha t})x_1x_3 + (C(e^{-\alpha t} - 1) + At^2e^{-\alpha t} + 2Bte^{-\alpha t})x_3^2, (x_1 + (e^{\alpha t} - 1)x_2 + tx_3)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\alpha \neq 0$,
- 27) $\varphi_t(x) = [e^{\delta t}x_0x_2 - 2Ate^{\delta t}x_1x_2 + Ate^{\delta t}x_2^2 - (2Bt + At^2)e^{\delta t}x_2x_3 + Ate^{\delta t}x_1^2 + (2Bt + At^2)e^{\delta t}x_1x_3 + (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t}x_3^2, (x_1 + (e^{-\delta t} - 1)x_2 + tx_3)x_2, e^{-\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- 28) $\varphi_t(x) = [e^{\delta t}x_0x_2 - 2A(e^{-\delta t} - e^{\delta t})x_1x_2 + (bte^{\delta t} + A(e^{-\delta t} - e^{\delta t}))x_2^2 + (-2Ate^{-\delta t} + 2B(1 - e^{-\delta t}))x_2x_3 + A(e^{-\delta t} - e^{\delta t})x_1^2 + 2(B(e^{-\delta t} - e^{\delta t}) + Ate^{-\delta t})x_1x_3 + (C(e^{-\delta t} - e^{\delta t}) + At^2e^{-\delta t} + 2Bte^{-\delta t})x_3^2, (x_1 + (e^{\delta t} - 1)x_2 + tx_3)x_2, e^{\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- 29) $\varphi_t(x) = [e^{\delta t}x_0x_3 + A(e^{2\alpha t} - e^{\delta t})x_1^2 + 2(B(e^{2\alpha t} - e^{\delta t}) + Ate^{2\alpha t})x_1x_2 + (C(e^{2\alpha t} - e^{\delta t}) + (At^2 + 2Bt)e^{2\alpha t})x_2^2, e^{\alpha t}(x_1 + tx_2)x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
- 30) $\varphi_t(x) = [x_0x_3 + ct x_3^3 + A(e^{2\alpha t} - 1)x_1^2 + 2(B(e^{2\alpha t} - 1) + Ate^{2\alpha t})x_1x_2 + (C(e^{2\alpha t} - 1) + (At^2 + 2Bt)e^{2\alpha t})x_2^2, e^{\alpha t}(x_1 + tx_2)x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,
- 31) $\varphi_t(x) = [e^{\delta t}x_0x_3 + ate^{\delta t}x_1x_3 + \frac{a}{2}t^2e^{\delta t}x_2x_3 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2(B(e^{2\delta t} - e^{\delta t}) + Ate^{2\delta t})x_1x_2 + (C(e^{2\delta t} - e^{\delta t}) + (At^2 + 2Bt)e^{2\delta t})x_2^2, e^{\delta t}(x_1 + tx_2)x_3, e^{\delta t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
- 32) $\varphi_t(x) = [e^{2\alpha t}x_0x_3 + Ate^{2\alpha t}x_1^2 + (2Bt + At^2)e^{2\alpha t}x_1x_2 + (Ct + Bt^2 + \frac{A}{3}t^3)e^{2\alpha t}x_2^2, e^{\alpha t}(x_1 + tx_2)x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\delta \neq 0$.
- 33) $\varphi_t(x) = [e^{\delta t}x_0x_2 + 2Ate^{\alpha t}x_1x_2 + At^2e^{\alpha t}x_2^2 + 2tBx_2x_3 + A(e^{\alpha t} - e^{\delta t})x_1^2 + 2B(1 - e^{\delta t})x_1x_3 + C(e^{-\alpha t} - e^{\delta t})x_3^2, e^{\alpha t}(x_1 + tx_2)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- 34) $\varphi_t(x) = [x_0x_2 + 2Ate^{\alpha t}x_1x_2 + At^2e^{\alpha t}x_2^2 + (ct + Bt^2)x_2x_3 + A(e^{\alpha t} - 1)x_1^2 + 2Btx_1x_3 + C(e^{-\alpha t} - 1)x_3^2, e^{\alpha t}(x_1 + tx_2)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\alpha \neq 0$,
- 35) $\varphi_t(x) = [e^{\delta t}x_0x_2 + (at + At^2)e^{\delta t}x_1x_2 + (\frac{a}{2}t^2 + \frac{A}{3}t^3)e^{\delta t}x_2^2 + 2tBx_2x_3 + Ate^{\delta t}x_1^2 + 2B(1 - e^{\delta t})x_1x_3 + C(e^{-\delta t} - e^{\delta t})x_3^2, e^{\delta t}(x_1 + tx_2)x_2, e^{\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- 36) $\varphi_t(x) = [e^{\delta t}x_0x_2 + 2Ate^{-\delta t}x_1x_2 + At^2e^{-\delta t}x_2^2 + 2Btx_2x_3 + A(e^{-\delta t} - e^{\delta t})x_1^2 + 2B(1 - e^{\delta t})x_1x_3 + Cte^{\delta t}x_3^2, e^{-\delta t}(x_1 + tx_2)x_2, e^{-\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$.
- 37) $\varphi_t(x) = [e^{\delta t}x_0x_3 + (At^2 + 2tB)x_1x_3 + (At^3 + 3Bt^2 + 2Ct)x_2x_3 + (\frac{A}{4}t^4 + Bt^3 + Ct^2)x_3^2 + A(1 - e^{\delta t})x_1^2 + 2(B(1 - e^{\delta t}) + At)x_1x_2 + (C(1 - e^{\delta t}) + 2Bt + At^2)x_2^2, (x_1 + tx_2 + \frac{t^2}{2}x_3)x_3, (x_2 + tx_3)x_3, x_3^2]$ with $\delta \neq 0$.

where all the parameters are constant complex numbers. Moreover, two such flows φ_t^1 and φ_t^2 are linearly conjugated if and only if they are of the same type with the same δ, α, β and there exists $\lambda \in \mathbb{C}^*$ such that $(A_1, B_1, C_1, a_1, b_1, c_1) = \lambda(A_2, B_2, C_2, a_2, b_2, c_2)$ unless φ_t^1 and φ_t^2 are both in the cases II11 or II14 and the linear conjugation switches x_1 and x_2 .

Remark 3.34. As in the previous cases up to a normalization of the time coordinate t we can assume that: in case **I**) either α, β or μ are 1 (if not vanishing) and in case **II**) that either $\delta = 1$ or we have $\delta = 0$ and α or β are 1 (if not vanishing).

Example 3.35. Note that in case II12 for $A = B = C = 1$ we have $C_{It} = \{((e^{\alpha t} - 1)x_1 + (e^{\beta t} - 1)x_2) \cdot ((e^{\alpha t} + 1)x_1 + (e^{\beta t} + 1)x_2) = x_3 = 0\}$. Therefore $\varphi_t \in \mathbf{osc}^{[2]}(\times)$ for all $t \neq 0$ and there is no line $L \subset C_{It}$ which is fix. However φ_t preserves the families of hyperplanes through $x_1 = x_3 = 0$ or $x_2 = x_3 = 0$.

Indeed we have:

Corollary 3.36. *Let φ_t be a quadratic flow in $\mathbf{tan}^{[2]}(//) \cup \mathbf{osc}^{[2]}(\times) \cup \mathbf{lin}$. Then there is (at least) a line L_I such that φ_t preserves the family of hyperplanes through L_I .*

Proof. It is enough to take $L_I = \{x_1 = x_3 = 0\}$ for flows in case **I**) and for flows II6 and II9 in case **II**) and $L_I = \{x_2 = x_3 = 0\}$ for the rest of flows. $\square$

Remark 3.37. Even in the case when P_I is fix one cannot obtain a result in the spirit of Corollary 3.22 for $P_I \in L_I$ for every possible choice of L_I as in the previous Corollary.

Proof. (of theorem 3.33) We have already seen that H is fix and that φ_t is a polynomial flow. We will start considering the case P_I mobile. We resume the notation and computations of lemma 3.31. As $\tilde{\varphi}_0 = \text{Id}$ we have $\alpha_0 = \gamma_0 = 0$ and $\tilde{\ell}_{0_0} = x$, $\tilde{\ell}_{1_0} = y$ and $\tilde{\ell}_{2_0} = z$. Let us impose now $\tilde{\varphi}_{t+s} = \tilde{\varphi}_s \circ \tilde{\varphi}_t$. Using again that P_t is mobile one concludes that $\tilde{\ell}_{1_t} = \mu_t \cdot y$ (note that otherwise the second component of the composition $\tilde{\varphi}_s \circ \tilde{\varphi}_t$ would have a quadratic term) and, as φ_t is a flow, $\mu_t = e^{\mu t}$. The first component of $\tilde{\varphi}_s \circ \tilde{\varphi}_t$ is

$$\alpha_s e^{2\mu t} y^2 + A_s^0 \alpha_t y^2 + A_s^0 \tilde{\ell}_{0_t} + B_s^0 e^{\mu t} \tilde{\ell}_{1_t} + C_s^0 \gamma_t y^2 + C_s^0 \tilde{\ell}_{2_t} + D_s^0 = (\alpha_s e^{2\mu t} + A_s^0 \alpha_t + C_s^0 \gamma_t) y^2 + \tilde{\ell}_{0_s}(\tilde{\ell}_{i_t}),$$

which yields

$$(3.27) \quad \alpha_{t+s} = e^{2\mu t} \alpha_s + A_s^0 \alpha_t + C_s^0 \gamma_t$$

and $\tilde{\ell}_{0_s}(\tilde{\ell}_{i_t}) = \tilde{\ell}_{0_{t+s}}$. Analogously, for the third component one obtains

$$(3.28) \quad \gamma_{t+s} = e^{2\mu t} \gamma_s + A_s^2 \alpha_t + C_s^2 \gamma_t$$

and $\tilde{\ell}_{2_s}(\tilde{\ell}_{i_t}) = \tilde{\ell}_{2_{t+s}}$. In particular $\tilde{\Psi}_t = (\tilde{\ell}_{0_t}, e^{\mu t} y, \tilde{\ell}_{2_t})$ is a linear flow in $\mathbb{C}^3$. A direct computation shows that there are the following possibilities for $\tilde{\Psi}_t$:

- (i) $\tilde{\Psi}_t = (e^{\alpha t} x, e^{\mu t} y, e^{\beta t} z)$,
- (ii) $\tilde{\Psi}_t = (e^{\mu t} x + t e^{\mu t} y, e^{\mu t} y, e^{\beta t} z)$,
- (iii) $\tilde{\Psi}_t = (e^{\alpha t} x, e^{\mu t} y, z + t)$,
- (iv) $\tilde{\Psi}_t = (e^{\mu t} x + t e^{\mu t} y, e^{\mu t} y, z + t)$.

Therefore one can assume $C_t^0 = A_t^2 = 0$ and equations 3.27 and 3.28 yield

$$\alpha_t = a(e^{2\mu t} - A_t^0), \quad \gamma_t = b(e^{2\mu t} - C_t^2)$$

and note that A_t^0 is equal to $e^{\alpha t}$ or $e^{\mu t}$ and γ_t is equal to $e^{\beta t}$ or 1. We can assume $A_t^0 = e^{\alpha t}$ and $\gamma_t = e^{\beta t}$, we obtain then the list of flows in **I**) and the statement in **vii**).

Let us tackle now the case P_I fix. Note that we can use the same arguments as in lemma 3.32 to conclude that one can assume $P_I = [1, 0, 0, 0]$ and

$$\varphi_t[x_0, x_1, x_2, x_3] = [\delta_t x_0 + \frac{q_t}{\ell_t}, \ell_{1_t}, \ell_{2_t}, \ell_{3_t}]$$

with $\ell_{1_t}, \ell_{2_t}, \ell_{3_t}, \ell_t \in A_1(x_1, x_2, x_3)$, $q_t \in A_2(x_1, x_2, x_3)$, $q_0 = 0$, $\delta_t = e^{\delta t}$ and $\ell_{i_0} = x_i$ for $i = 1, 2, 3$. Moreover as $\Psi_t(x) = [\ell_{1_t}, \ell_{2_t}, \ell_{3_t}]$ is a linear flow on $\mathbb{P}^2$ it belongs to the following list:

- (i) $\Psi_t(x) = [e^{\alpha t} x_1, x_2 + t x_3, x_3]$,
- (ii) $\Psi_t(x) = [e^{\alpha t} x_1, e^{\beta t} x_2, x_3]$,
- (iii) $\Psi_t(x) = [x_1 + (e^{\alpha t} - 1)x_2 + t x_3, e^{\alpha t} x_2, x_3]$,
- (iv) $\Psi_t(x) = [e^{\alpha t}(x_1 + t x_2), e^{\alpha t} x_2, x_3]$,
- (v) $\Psi_t(x) = [x_1 + t x_2 + \frac{t^2}{2} x_3, x_2 + t x_3, x_3]$.

The statements in **v**) and **vi**) follow immediately. The condition $\varphi_{t+s} = \varphi_s \circ \varphi_t$ translates into

$$\ell(\ell_{i_t}) \cdot q_{t+s} = e^{\delta s} \cdot q_t \cdot \ell(\ell_{i_t}) + \ell \cdot q_s(\ell_{i_t})$$

or equivalently

$$(3.29) \quad \ell(\ell_{i_t}) \cdot (q_{t+s} - e^{\delta s} q_t) = \ell \cdot q_s(\ell_{i_t}).$$

If $(\ell(\ell_{i_t})) = 0$ were mobile then $q_{t+s} - e^{\delta s} q_t$ has ℓ as a factor, which implies (taking $t = 0$) that q_s has ℓ as a fixed factor. However this would yield φ_t linear for every t , which is a contradiction. Therefore

$$(3.30) \quad \ell(\ell_{i_t}) = \lambda(t) \ell$$

and

$$(3.31) \quad \lambda(t)(q_{t+s} - e^{\delta s} q_t) = q_s(\ell_{it}).$$

When we impose the condition 3.30 we conclude that we have the following possibilities:

- 1) $\ell = x_3$, $\lambda(t) = 1$ and Ψ_t is either of the previous flows,
- 2) $\ell = x_2$ and Ψ_t is of types **iii**) or **iv**) and $\lambda(t) = e^{\alpha t}$,
- 3) or $\ell = x_1$, $\lambda(t) = e^{\alpha t}$ and Ψ_t is of type **i**).

Indeed, notice that in case **ii**) up to a change of coordinates one can assume that $\ell = x_3$, which is the reason why we have excluded the other possibilities of the previous list. We proceed now to determine q_t in each case.

- 1) $\ell = x_3$, $\lambda(t) = 1$ and Ψ_t of the following type:

- i) We can assume $q_t(x) = x_3 v_t + \tilde{q}_t$ where $\tilde{q}_t \in A_2(x_1, x_2)$ and $v_t \in A_1(x_1, x_2, x_3)$. Equation 3.31 is rewritten as

$$(3.32) \quad x_3(v_{t+s} - e^{\delta s} v_t) + \tilde{q}_{t+s} - e^{\delta s} \tilde{q}_t = x_3 v_s(\ell_{it}) + \tilde{q}_s(\ell_{it}).$$

We denote $\tilde{q}_t = A(t)x_1^2 + 2B(t)x_1x_2 + C(t)x_2^2$ and $v_t = a(t)x_1 + b(t)x_2 + c(t)x_3$. Then equation 3.32 is equivalent to

$$\begin{aligned} A(t+s) - e^{\delta s} A(t) &= e^{2\alpha t} A(s) \\ B(t+s) - e^{\delta s} B(t) &= e^{\alpha t} B(s) \\ C(t+s) - e^{\delta s} C(t) &= C(s) \\ a(t+s) - e^{\delta s} a(t) &= e^{\alpha t} a(s) + 2te^{\alpha t} B(s) \\ b(t+s) - e^{\delta s} b(t) &= b(s) + 2tC(s) \\ c(t+s) - e^{\delta s} c(t) &= c(s) + t^2 C(s) + tb(s) \end{aligned}$$

yielding solutions

$$\begin{aligned} A(t) &= A(e^{2\alpha t} - e^{\delta t}) \quad (\text{or } Ate^{\delta t} \quad \text{for } \delta = 2\alpha) \\ B(t) &= B(e^{\alpha t} - e^{\delta t}) \quad (\text{or } Bte^{\delta t} \quad \text{for } \delta = \alpha) \\ C(t) &= C(1 - e^{\delta t}) \quad (\text{or } Ct \quad \text{for } \delta = 0) \\ a(t) &= a(e^{\alpha t} - e^{\delta t}) + 2te^{\alpha t} B \quad (\text{or } ate^{\delta t} + Bt^2 e^{\delta t} \quad \text{for } \delta = \alpha) \\ b(t) &= b(1 - e^{\delta t}) + 2tC \quad (\text{or } bt + Ct^2 \quad \text{for } \delta = 0) \\ c(t) &= c(1 - e^{\delta t}) + tb + t^2 C \quad (\text{or } ct + \frac{b}{2}t^2 + Ct^3 \quad \text{for } \delta = 0) \end{aligned}$$

and the flows

- $\varphi_t(x) = [e^{\delta t} x_0 x_3 + (a(e^{\alpha t} - e^{\delta t}) + 2tBe^{\alpha t}) x_1 x_3 + (b(1 - e^{\delta t}) + 2tC) x_2 x_3 + (c(1 - e^{\delta t}) + t^2 C + tb) x_3^2 + A(e^{2\alpha t} - e^{\delta t}) x_1^2 + 2B(e^{\alpha t} - e^{\delta t}) x_1 x_2 + C(1 - e^{\delta t}) x_2^2, e^{\alpha t} x_1 x_3, x_2 x_3 + tx_3^2, x_3^2]$ with $\delta \neq 0$, $\delta \neq \alpha$, $\delta \neq 2\alpha$.
- $\varphi_t(x) = [x_0 x_3 + (a(e^{\alpha t} - 1) + 2tBe^{\alpha t}) x_1 x_3 + (bt + Ct^2) x_2 x_3 + (ct + \frac{b}{2}t^2 + \frac{C}{3}t^3) x_3^2 + A(e^{2\alpha t} - 1) x_1^2 + 2B(e^{\alpha t} - 1) x_1 x_2 + Ct x_2^2, e^{\alpha t} x_1 x_3, x_2 x_3 + tx_3^2, x_3^2]$ with $\alpha \neq 0$,
- $\varphi_t(x) = [x_0 x_3 + (at + Bt^2) x_1 x_3 + (bt + Ct^2) x_2 x_3 + (ct + \frac{b}{2}t^2 + \frac{C}{3}t^3) x_3^2 + At x_1^2 + 2Bt x_1 x_2 + Ct x_2^2, x_1 x_3, x_2 x_3 + tx_3^2, x_3^2]$,
- $\varphi_t(x) = [e^{\delta t} x_0 x_3 + (at + Bt^2) e^{\delta t} x_1 x_3 + (b(1 - e^{\delta t}) + 2tC) x_2 x_3 + (c(1 - e^{\delta t}) + t^2 C + tb) x_3^2 + A(e^{2\delta t} - e^{\delta t}) x_1^2 + 2Bte^{\delta t} x_1 x_2 + C(1 - e^{\delta t}) x_2^2, e^{\delta t} x_1 x_3, x_2 x_3 + tx_3^2, x_3^2]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{\delta t} x_0 x_3 + (a(e^{\frac{\alpha t}{2}} - e^{\delta t}) + 2tBe^{\frac{\alpha t}{2}}) x_1 x_3 + (b(1 - e^{\delta t}) + 2tC) x_2 x_3 + (c(1 - e^{\delta t}) + t^2 C + tb) x_3^2 + Ate^{\delta t} x_1^2 + 2B(e^{\frac{\alpha t}{2}} - e^{\delta t}) x_1 x_2 + C(1 - e^{\delta t}) x_2^2, e^{\frac{\alpha t}{2}} x_1 x_3, x_2 x_3 + tx_3^2, x_3^2]$ with $\delta \neq 0$.

For the rest of the cases see the analogous computations in the Appendix A.

We will discuss now when two flows of the previous list are linearly conjugated. Let φ_t^1 and φ_t^2 be two flows of the previous lists such that there is a linear conjugation $A = [\mu_0, \mu_1, \mu_2, \mu_3]$ with $\mu_i \in A_1(x_0, x_1, x_2, x_3)$ such that

$$\varphi_t^2 = A^{-1} \circ \varphi_t^1 \circ A.$$

It is not difficult to convince oneself that in order to be linearly conjugated φ_t^1 and φ_t^2 must be both of type **I)** or **II)**.

Assume that they are of type **I)**. As $H = (x_3 = 0)$ is fix we can assume that $\mu_3 = x_3$ and that A induces a linear conjugation between the induced polynomial flows $\tilde{\varphi}_t^1$ and $\tilde{\varphi}_t^2$ in $\mathbb{C}^3$. One can check that the only possibilities of conjugation are between flows of the same type (**a**), **b**), **c**) or **d**)) with $\alpha_1 = \alpha_2$ and $\beta_1 = \beta_2$. In this case the only conjugations are those which allow to assume that the pair (a, b) is of the form $(a, 1)$ with $a \in \mathbb{C}^*$.

We assume now that φ_t^1, φ_t^2 are two flows of type **II)** and $A = [\mu_0, \mu_1, \mu_2, \mu_3]$ with $\mu_i \in A_1(x_0, x_1, x_2, x_3)$ such that

$$\varphi_t^2 = A^{-1} \circ \varphi_t^1 \circ A.$$

As $P_I = [1, 0, 0, 0]$ and the planes H^1 and H^2 corresponding to the flows φ_t^1 and φ_t^2 respectively are of the form $H^1 = (x_{i_1} = 0)$ or $H^2 = (x_{i_2} = 0)$, we have $\mu_i \in A_i(x_1, x_2, x_3)$ for $i = 1, 2, 3$ and $\mu_{i_2} = \mu_{i_2}^{i_1} x_{i_1}$. Moreover we can choose $\mu_{i_2}^{i_1} = 1$. This implies that the linear flows in Ψ_t^1, Ψ_t^2 in $\mathbb{P}^2$ corresponding to φ_t^1, φ_t^2 (same notation as before) are linearly conjugated by $\tilde{A} = [\mu_1, \mu_2, \mu_3]$. One concludes that $\tilde{A} = [x_1, x_2, x_3]$ and $\Psi_t^1 = \Psi_t^2$. In particular $i_1 = i_2$ so $H^1 = H^2$. We denote now

$$\mu_0(x) = \mu_0^0 x_0 + \mu_0^1 x_1 + \mu_0^2 x_2 + \mu_0^3 x_3$$

and impose $A \circ \varphi_t^2 = \varphi_t^1 \circ A$ on the first component for each of the previous cases. Let us make the computations for the first case (which corresponds to the flow 1):

$$\begin{aligned} & \mu_0^0 (e^{\delta_2 t} x_0 x_3 + (a_2(e^{\alpha_2 t} - e^{\delta_2 t}) + 2tB_2 e^{\alpha_2 t}) x_1 x_3 + (b_2(1 - e^{\delta_2 t}) + 2tC_2) x_2 x_3 + (c_2(1 - e^{\delta_2 t}) + t^2 C_2 + tb_2) x_3^2 + \\ & A_2(e^{2\alpha_2 t} - e^{\delta_2 t}) x_1^2 + 2B_2(e^{\alpha_2 t} - e^{\delta_2 t}) x_1 x_2 + C_2(1 - e^{\delta_2 t}) x_2^2) + \mu_0^1 e^{\alpha_2 t} x_1 x_3 + \mu_0^2 x_2 x_3 + \mu_0^3 x_3^2 = \\ & e^{\delta_1 t} (\mu_0^0 x_0 + \mu_0^1 x_1 + \mu_0^2 x_2 + \mu_0^3 x_3) + (a_1(e^{\alpha_1 t} - e^{\delta_1 t}) + 2tB_1 e^{\alpha_1 t}) x_1 x_3 + (b_1(1 - e^{\delta_1 t}) + 2tC_1) x_2 x_3 + \\ & (c_1(1 - e^{\delta_1 t}) + t^2 C_1 + tb_1) x_3^2 + A_1(e^{2\alpha_1 t} - e^{\delta_1 t}) x_1^2 + 2B_1(e^{\alpha_1 t} - e^{\delta_1 t}) x_1 x_2 + C_1(1 - e^{\delta_1 t}) x_2^2. \end{aligned}$$

It follows that $\delta_1 = \delta_2$ and

$$\begin{aligned} A_1 &= \mu_0^0 A_2 \\ B_1 &= \mu_0^0 B_2 \\ C_1 &= \mu_0^0 C_2 \\ a_1 &= \mu_0^0 a_2 + \mu_0^1 \\ b_1 &= \mu_0^0 b_2 + \mu_0^2 \\ c_1 &= \mu_0^0 c_2 + \mu_0^3. \end{aligned}$$

As $\mu_0^0 \in \mathbb{C}^*$ and $\mu_0^1, \mu_0^2, \mu_0^3 \in \mathbb{C}$ we can assume that $a_1 = b_1 = c_1 = 0$ and A_1, B_1, C_1 are defined modulo a $\mathbb{C}^*$ action. Therefore we obtain the flow in II1 and we conclude that two flows of this type are linearly conjugated if and only if there exists $\mu_0^0 \in \mathbb{C}^*$ such that $(A_1, B_1, C_1) = \mu_0^0 (A_2, B_2, C_2)$. Analogous calculations yield the flows II2, II3, II4 and II5.

Analogously we obtain the following flows for each of the cases: **1) ii)** from II11 to II20, **1) iii)** from II21 to II24, **1) iv)** from II29 to II32, **1) v)** II37, **2) iii)** from II25 to II28, **2) iv)** from II33 to II36 and **3) iii)** from II6 to II10. Note that in some cases due to simmetry some extra considerations are necessary. $\square$

3.4. Conclusions. We can sum up the results of this section in the following results:

Theorem 3.38. *Let φ_t be a quadratic flow in Bir^3 . Then one of the following possibilities hold:*

- a)** $\varphi_t \in \text{gen}^{[2]}(//) \cup \text{lin}$, P_I, S, C_I are fix and H_t is mobile,
- b)** $\varphi_t \in \text{tan}^{[2]}(\text{O}) \cup \text{tan}^{[2]}(\times) \cup \text{lin}$, P_I, S are fix and H_t, C_{I_t} are mobile,
- c)** $\varphi_t \in \text{tan}^{[2]}(//) \cup \text{lin}$, H, C_I are fix and P_{I_t} can be either fix or mobile,
- d)** $\varphi_t \in \text{osc}^{[2]}(\times) \cup \text{lin}$, H, P_I are fix and C_I can be either fix or mobile.

Moreover, if H is fix then φ_t is a polynomial flow, i.e. $\varphi_t|_{\mathbb{P}^3 \setminus H} : \mathbb{P}^3 \setminus H \rightarrow \mathbb{P}^3 \setminus H$ is polynomial for each t . In particular there are no flows in $\text{gen}^{[2]}(\text{O}) \cup \text{gen}^{[2]}(\times) \cup \text{lin}$.

Theorem 3.39. *Let φ_t be a quadratic flow in $\mathbf{Bir}^3$. Then:*

- i) *There exists a line L such that φ_t preserves the family of hyperplanes through L . Moreover in cases a) and c) we can choose $L = C_I$.*
- ii) *If P_I is fix then φ_t preserves the family of hyperplanes through P_I (in particular the family of lines through P_I).*

Corollary 3.40. *Let φ_t a quadratic generic flow in $\mathbf{Bir}^3$. Then φ_t is determined by a linear flow η_t on the $\mathbb{P}^2$ of the net of lines through P_I and a linear flow χ_t on the $\mathbb{P}^1$ of the pencil of planes through the line C_I . Namely*

$$\varphi_t(P) = \eta_t(P \vee P_I) \cap \chi_t(P \vee C_I).$$

Theorem 3.41. *Let φ_t be a quadratic flow in $\mathbf{Bir}^3$. Then, up to linear conjugation, φ_t is in one (and only one) of the lists of theorems 3.20 (generic case), 3.26 (non-generic with conic C_I of multiplicity 1 at P_I) and 3.33 (non-generic with conic C_I of multiplicity 2 at P_I).*

Remark 3.42. As in $\mathbb{P}^2$ using the classification one verifies that if φ_t is a germ of quadratic flow then φ_t is defined for every $t \in \mathbb{C}$.

APPENDIX A. THE COMPUTATIONS OF THEOREM 3.26

We resume here the computations where we left them, using the same notation.

- 1) $\ell = x_3$, $\lambda(t) = 1$ and Ψ_t of the following type:

- ii) One obtains the equations

$$\begin{aligned} A(t+s) - e^{\delta s} A(t) &= e^{2\alpha t} A(s) \\ B(t+s) - e^{\delta s} B(t) &= e^{\alpha t} e^{\beta t} B(s) \\ C(t+s) - e^{\delta s} C(t) &= e^{2\beta t} C(s) \\ a(t+s) - e^{\delta s} a(t) &= e^{\alpha t} a(s) \\ b(t+s) - e^{\delta s} b(t) &= e^{\beta t} b(s) \\ c(t+s) - e^{\delta s} c(t) &= c(s) \end{aligned}$$

and the solutions

$$\begin{aligned} A(t) &= A(e^{2\alpha t} - e^{\delta t}) \quad (\text{or } Ate^{\delta t} \text{ for } \delta = 2\alpha) \\ B(t) &= B(e^{(\alpha+\beta)t} - e^{\delta t}) \quad (\text{or } Bte^{\delta t} \text{ for } \delta = \alpha + \beta) \\ C(t) &= C(e^{2\beta t} - e^{\delta t}) \quad (\text{or } Cte^{\delta t} \text{ for } \delta = 2\beta) \\ a(t) &= a(e^{\alpha t} - e^{\delta t}) \quad (\text{or } ate^{\delta t} \text{ for } \delta = \alpha) \\ b(t) &= b(e^{\beta t} - e^{\delta t}) \quad (\text{or } bte^{\delta t} \text{ for } \delta = \beta) \\ c(t) &= c(1 - e^{\delta t}) \quad (\text{or } ct \text{ for } \delta = 0) \end{aligned}$$

Up to a linear conjugation switching x_1 and x_2 we obtain the following list of flows from the previous solutions:

- $\varphi_t(x) = [e^{\delta t} x_0 x_3 + a(e^{\alpha t} - e^{\delta t}) x_1 x_3 + b(e^{\beta t} - e^{\delta t}) x_2 x_3 + c(1 - e^{\delta t}) x_3^2 + A(e^{2\alpha t} - e^{\delta t}) x_1^2 + 2B(e^{(\alpha+\beta)t} - e^{\delta t}) x_1 x_2 + C(e^{2\beta t} - e^{\delta t}) x_2^2, e^{\alpha t} x_1 x_3, e^{\beta t} x_2 x_3, x_3^2]$ with $\delta \neq 0$, $\delta \neq \alpha, \beta, \alpha + \beta, 2\alpha, 2\beta$,
- $\varphi_t(x) = [x_0 x_3 + a(e^{\alpha t} - 1) x_1 x_3 + b(e^{\beta t} - 1) x_2 x_3 + ct x_3^2 + A(e^{2\alpha t} - 1) x_1^2 + 2B(e^{(\alpha+\beta)t} - 1) x_1 x_2 + C(e^{2\beta t} - 1) x_2^2, e^{\alpha t} x_1 x_3, e^{\beta t} x_2 x_3, x_3^2]$ with $\alpha \neq 0$ and $\beta \neq 0$,
- $\varphi_t(x) = [e^{\delta t} x_0 x_3 + ate^{\delta t} x_1 x_3 + b(e^{\beta t} - e^{\delta t}) x_2 x_3 + c(1 - e^{\delta t}) x_3^2 + A(e^{2\delta t} - e^{\delta t}) x_1^2 + 2B(e^{(\delta+\beta)t} - e^{\delta t}) x_1 x_2 + C(e^{2\beta t} - e^{\delta t}) x_2^2, e^{\delta t} x_1 x_3, e^{\beta t} x_2 x_3, x_3^2]$ with $\delta \neq 0$, $\delta = \alpha$, $\delta \neq \beta, 2\beta$,
- $\varphi_t(x) = [e^{2\alpha t} x_0 x_3 + a(e^{\alpha t} - e^{2\alpha t}) x_1 x_3 + b(e^{\beta t} - e^{2\alpha t}) x_2 x_3 + c(1 - e^{2\alpha t}) x_3^2 + Ate^{2\alpha t} x_1^2 + 2B(e^{(\alpha+\beta)t} - e^{2\alpha t}) x_1 x_2 + C(e^{2\beta t} - e^{2\alpha t}) x_2^2, e^{\alpha t} x_1 x_3, e^{\beta t} x_2 x_3, x_3^2]$ with $\alpha \neq 0$, $\beta \neq 0$, $\alpha \neq \beta$,
- $\varphi_t(x) = [e^{\delta t} x_0 x_3 + ate^{\delta t} x_1 x_3 + bte^{\delta t} x_2 x_3 + c(1 - e^{\delta t}) x_3^2 + A(e^{2\delta t} - e^{\delta t}) x_1^2 + 2B(e^{2\delta t} - e^{\delta t}) x_1 x_2 + C(e^{2\delta t} - e^{\delta t}) x_2^2, e^{\delta t} x_1 x_3, e^{\delta t} x_2 x_3, x_3^2]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{2\alpha t} x_0 x_3 + a(e^{\alpha t} - e^{2\alpha t}) x_1 x_3 + b(e^{\alpha t} - e^{2\alpha t}) x_2 x_3 + c(1 - e^{2\alpha t}) x_3^2 + Ate^{2\alpha t} x_1^2 + 2Bte^{2\alpha t} x_1 x_2 + Cte^{2\alpha t} x_2^2, e^{\alpha t} x_1 x_3, e^{\alpha t} x_2 x_3, x_3^2]$ with $\alpha \neq 0$,

- $\varphi_t(x) = [e^{(\alpha+\beta)t}x_0x_3 + a(e^{\alpha t} - e^{(\alpha+\beta)t})x_1x_3 + b(e^{\beta t} - e^{(\alpha+\beta)t})x_2x_3 + c(1 - e^{(\alpha+\beta)t})x_3^2 + A(e^{2\alpha t} - e^{(\alpha+\beta)t})x_1^2 + 2Bte^{(\alpha+\beta)t}x_1x_2 + C(e^{2\beta t} - e^{(\alpha+\beta)t})x_2^2, e^{\alpha t}x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\alpha \neq 0, \beta \neq 0, \alpha \neq \beta$,
- $\varphi_t(x) = [e^{\delta t}x_0x_3 + ate^{\delta t}x_1x_3 + b(1 - e^{\delta t})x_2x_3 + c(1 - e^{\delta t})x_3^2 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2Bte^{\delta t}x_1x_2 + C(1 - e^{\delta t})x_2^2, e^{\delta t}x_1x_3, x_2x_3, x_3^2]$ with $\delta \neq 0$,
- $\varphi_t(x) = [x_0x_3 + atx_1x_3 + b(e^{\beta t} - 1)x_2x_3 + ct x_3^2 + Atx_1^2 + 2B(e^{\beta t} - 1)x_1x_2 + C(e^{2\beta t} - 1)x_2^2, x_1x_3, e^{\beta t}x_2x_3, x_3^2]$ with $\beta \neq 0$,
- $\varphi_t(x) = [x_0x_3 + t((ax_1 + bx_2 + cx_3)x_3 + (Ax_1^2 + 2Bx_1x_2 + Cx_2^2)), x_1x_3, x_2x_3, x_3^2]$,

iii) Again with the same notation we obtain:

$$\begin{aligned}
 A(t+s) - e^{\delta s}A(t) &= A(s) \\
 B(t+s) - e^{\delta s}B(t) &= e^{\alpha t}B(s) + A(s)(e^{\alpha t} - 1) \\
 C(t+s) - e^{\delta s}C(t) &= C(s)e^{2\alpha t} + (e^{\alpha t} - 1)^2A(s) + 2B(s)e^{\alpha t}(e^{\alpha t} - 1) \\
 a(t+s) - e^{\delta s}a(t) &= a(s) + 2tA(s) \\
 b(t+s) - e^{\delta s}b(t) &= e^{\alpha t}b(s) + (e^{\alpha t} - 1)a(s) + 2t(e^{\alpha t} - 1)A(s) + 2te^{\alpha t}B(s) \\
 c(t+s) - e^{\delta s}c(t) &= c(s) + t^2A(s) + ta(s)
 \end{aligned}$$

yielding solutions

$$\begin{aligned}
 A(t) &= A(1 - e^{\delta t}) \quad (\text{or } At \text{ for } \delta = 0) \\
 B(t) &= B(e^{\alpha t} - e^{\delta t}) - A(1 - e^{\delta t}) \\
 &\quad (\text{or } B(e^{\delta t} - 1) - At \text{ for } \delta = 0, \alpha \neq 0, \\
 &\quad \text{or } Bte^{\delta t} - A(1 - e^{\delta t}) \text{ for } \alpha = \delta \neq 0), \\
 C(t) &= C(e^{2\alpha t} - e^{\delta t}) - 2B(e^{\alpha t} - e^{\delta t}) + A(1 - e^{\delta t}) \\
 &\quad (\text{or } C(e^{2\alpha t} - 1) - 2B(e^{\alpha t} - 1) + At \text{ for } \delta = 0, \alpha \neq 0, \\
 &\quad \text{or } C(e^{2\delta t} - e^{\delta t}) - 2Bte^{\delta t} + A(1 - e^{\delta t}) \text{ for } \delta = \alpha \neq 0, \\
 &\quad \text{or } Cte^{2\alpha t} - 2B(e^{\alpha t} - e^{2\alpha t}) + A(1 - e^{2\alpha t}) \text{ for } \delta = 2\alpha, \alpha \neq 0), \\
 a(t) &= a(1 - e^{\delta t}) + 2tA \quad (\text{or } at + At^2 \text{ for } \delta = 0) \\
 b(t) &= b(e^{\alpha t} - e^{\delta t}) + 2te^{\alpha t}B - a(1 - e^{\delta t}) - 2tA \\
 &\quad (\text{or } b(e^{\alpha t} - 1) + 2te^{\alpha t}B - At^2 - at \text{ for } \delta = 0, \alpha \neq 0, \\
 &\quad \text{or } (Bt^2 + bt + a)e^{\delta t} - 2At - a \text{ for } \alpha = \delta \neq 0), \\
 c(t) &= c(1 - e^{\delta t}) + at + At^2 \quad (\text{or } ct + \frac{a}{2}t^2 + \frac{A}{3}t^3 \text{ for } \delta = 0).
 \end{aligned}$$

Note that we can assume $\alpha \neq 0$ (otherwise we obtain one of the flows in **i**) and that the equations obtained for $A(t) + B(t), C(t) + 2B(t) + A(t)$ and $a(t) + b(t)$ are easy to solve. We obtain the flows:

- $\varphi_t(x) = [e^{\delta t}x_0x_3 + (a(1 - e^{\delta t}) + 2tA)x_1x_3 + (b(e^{\alpha t} - e^{\delta t}) + 2te^{\alpha t}B - a(1 - e^{\delta t}) - 2tA)x_2x_3 + (c(1 - e^{\delta t}) + at + At^2)x_3^2 + A(1 - e^{\delta t})x_1^2 + 2(B(e^{\alpha t} - e^{\delta t}) - A(1 - e^{\delta t}))x_1x_2 + (C(e^{2\alpha t} - e^{\delta t}) - 2B(e^{\alpha t} - e^{\delta t}) + A(1 - e^{\delta t}))x_2^2, x_1x_3 + (e^{\alpha t} - 1)x_2x_3 + tx_3^2, e^{\alpha t}x_2x_3, x_3^2]$ with $\delta \neq 0, \alpha, 2\alpha, \alpha \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_3 + (a(1 - e^{\delta t}) + 2tA)x_1x_3 + ((Bt^2 + bt + a)e^{\delta t} - 2At - a)x_2x_3 + (c(1 - e^{\delta t}) + at + At^2)x_3^2 + A(1 - e^{\delta t})x_1^2 + 2(Bte^{\delta t} - A(1 - e^{\delta t}))x_1x_2 + (C(e^{2\delta t} - e^{\delta t}) - 2Bte^{\delta t} + A(1 - e^{\delta t}))x_2^2, x_1x_3 + (e^{\delta t} - 1)x_2x_3 + tx_3^2, e^{\delta t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{2\alpha t}x_0x_3 + (a(1 - e^{2\alpha t}) + 2tA)x_1x_3 + (b(e^{\alpha t} - e^{2\alpha t}) + 2te^{\alpha t}B - a(1 - e^{2\alpha t}) - 2tA)x_2x_3 + (c(1 - e^{2\alpha t}) + at + At^2)x_3^2 + A(1 - e^{2\alpha t})x_1^2 + 2(B(e^{\alpha t} - e^{2\alpha t}) - A(1 - e^{2\alpha t}))x_1x_2 + (Cte^{2\alpha t} - 2B(e^{\alpha t} - e^{2\alpha t}) + A(1 - e^{2\alpha t}))x_2^2, x_1x_3 + (e^{\alpha t} - 1)x_2x_3 + tx_3^2, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,
- $\varphi_t(x) = [x_0x_3 + (at + At^2)x_1x_3 + (b(e^{\alpha t} - 1) + 2Bte^{\alpha t} - At^2 - at)x_2x_3 + (\frac{A}{3}t^3 + \frac{a}{2}t^2 + ct)x_3^2 + Atx_1^2 + 2(B(e^{\alpha t} - 1) - At)x_1x_2 + (C(e^{2\alpha t} - 1) - 2B(e^{\alpha t} - 1) + At)x_2^2, x_1x_3 + (e^{\alpha t} - 1)x_2x_3 + tx_3^2, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,

iv) One has the equations:

$$\begin{aligned}
A(t+s) - e^{\delta s}A(t) &= e^{2\alpha t}A(s) \\
B(t+s) - e^{\delta s}B(t) &= e^{2\alpha t}B(s) + te^{2\alpha t}A(s) \\
C(t+s) - e^{\delta s}C(t) &= e^{2\alpha t}(C(s) + 2tB(s) + t^2A(s)) \\
a(t+s) - e^{\delta s}a(t) &= e^{\alpha t}a(s) \\
b(t+s) - e^{\delta s}b(t) &= e^{\alpha t}b(s) + te^{\alpha t}a(s) \\
c(t+s) - e^{\delta s}c(t) &= c(s)
\end{aligned}$$

yielding solutions

$$\begin{aligned}
A(t) &= A(e^{2\alpha t} - e^{\delta t}) \quad (\text{or } Ate^{\delta t} \text{ for } \delta = 2\alpha) \\
B(t) &= B(e^{2\alpha t} - e^{\delta t}) + Ate^{2\alpha t} \quad (\text{or } (Bt + \frac{A}{2}t^2)e^{\delta t} \text{ for } \delta = 2\alpha) \\
C(t) &= C(e^{2\alpha t} - e^{\delta t}) + (At^2 + 2tB)e^{2\alpha t} \quad (\text{or } (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t} \text{ for } \delta = 2\alpha) \\
a(t) &= a(e^{\alpha t} - e^{\delta t}) \quad (\text{or } ate^{\delta t} \text{ for } \delta = \alpha) \\
b(t) &= b(e^{\alpha t} - e^{\delta t}) + ate^{\alpha t} \quad (\text{or } (bt + \frac{a}{2}t^2)e^{\delta t} \text{ for } \delta = \alpha) \\
c(t) &= c(1 - e^{\delta t}) \quad (\text{or } ct \text{ for } \delta = 0).
\end{aligned}$$

As before we can assume $\alpha \neq 0$. We obtain the flows:

- $\varphi_t(x) = [e^{\delta t}x_0x_3 + a(e^{\alpha t} - e^{\delta t})x_1x_3 + (b(e^{\alpha t} - e^{\delta t}) + ate^{\delta t})x_2x_3 + c(1 - e^{\delta t})x_3^2 + A(e^{2\alpha t} - e^{\delta t})x_1^2 + 2(B(e^{2\alpha t} - e^{\delta t}) + Ate^{2\alpha t})x_1x_2 + (C(e^{2\alpha t} - e^{\delta t}) + (At^2 + 2Bt)e^{2\alpha t})x_2^2, e^{\alpha t}(x_1 + tx_2)x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
- $\varphi_t(x) = [x_0x_3 + a(e^{\alpha t} - 1)x_1x_3 + (b(e^{\alpha t} - 1) + at)x_2x_3 + ct x_3^3 + A(e^{2\alpha t} - 1)x_1^2 + 2(B(e^{2\alpha t} - 1) + Ate^{2\alpha t})x_1x_2 + (C(e^{2\alpha t} - 1) + (At^2 + 2Bt)e^{2\alpha t})x_2^2, e^{\alpha t}(x_1 + tx_2)x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\alpha \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_3 + ate^{\delta t}x_1x_3 + (bt + a\frac{t^2}{2})e^{\delta t}x_2x_3 + c(1 - e^{\delta t})x_3^2 + A(e^{2\delta t} - e^{\delta t})x_1^2 + 2(B(e^{2\delta t} - e^{\delta t}) + Ate^{2\delta t})x_1x_2 + (C(e^{2\delta t} - e^{\delta t}) + (At^2 + 2Bt)e^{2\delta t})x_2^2, e^{\delta t}(x_1 + tx_2)x_3, e^{\delta t}x_2x_3, x_3^2]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{2\alpha t}x_0x_3 + a(e^{\alpha t} - e^{2\alpha t})x_1x_3 + (b(e^{\alpha t} - e^{2\alpha t}) + ate^{2\alpha t})x_2x_3 + c(1 - e^{2\alpha t})x_3^2 + Ate^{2\alpha t}x_1^2 + 2(Bt + A\frac{t^2}{2})e^{2\alpha t}x_1x_2 + (Ct + Bt^2 + A\frac{t^3}{3})e^{2\alpha t}x_2^2, e^{\alpha t}(x_1 + tx_2)x_3, e^{\alpha t}x_2x_3, x_3^2]$ with $\delta \neq 0$.

v) Finally one has:

$$\begin{aligned}
A(t+s) - e^{\delta s}A(t) &= A(s) \\
B(t+s) - e^{\delta s}B(t) &= B(s) + tA(s) \\
C(t+s) - e^{\delta s}C(t) &= C(s) + 2tB(s) + t^2A(s) \\
a(t+s) - e^{\delta s}a(t) &= a(s) + t^2A(s) + 2tB(s) \\
b(t+s) - e^{\delta s}b(t) &= b(s) + ta(s) + t^3A(s) + 3t^2B(s) + 2tC(s) \\
c(t+s) - e^{\delta s}c(t) &= c(s) + tb(s) + \frac{t^2}{2}a(s) + \frac{t^4}{4}A(s) + t^3B(s) + t^2C(s).
\end{aligned}$$

One verifies that if $\delta = 0$ the only solutions correspond to linear flows, therefore we can assume $\delta \neq 0$ and we obtain the following solutions:

$$\begin{aligned}
A(t) &= A(1 - e^{\delta t}) \\
B(t) &= B(1 - e^{\delta t}) + At \\
C(t) &= C(1 - e^{\delta t}) - 2tB + At^2 \\
a(t) &= a(1 - e^{\delta t}) + At^2 + 2tB \\
b(t) &= b(1 - e^{\delta t}) + at + At^3 + 3Bt^2 + 2Ct \\
c(t) &= c(1 - e^{\delta t}) + bt + (\frac{a}{2} + C)t^2 + Bt^3 + \frac{A}{4}t^4,
\end{aligned}$$

which yield the flow

$$\begin{aligned} \varphi_t(x) = & [e^{\delta t}x_0x_3 + (a(1 - e^{\delta t}) + At^2 + 2tB)x_1x_2 + (b(1 - e^{\delta t}) + at + At^3 + 3Bt^2 + 2Ct)x_2x_3 + \\ & (c(1 - e^{\delta t}) + bt + \frac{a}{2}t^2 + \frac{A}{4}t^4 + Bt^3 + Ct^2)x_3^2 + A(1 - e^{\delta t})x_1^2 + 2(B(1 - e^{\delta t}) + At)x_1x_2 + \\ & (C(1 - e^{\delta t}) + 2Bt + At^2)x_2^2, (x_1 + tx_2 + \frac{t^2}{2}x_3)x_3, (x_2 + tx_3)x_3, x_3^2] \end{aligned}$$

with $\delta \neq 0$.

2) $\ell = x_2$ and Ψ_t of the following type:

iii) Then $\lambda(t) = e^{\alpha t}$ and we can assume $\alpha \neq 0$. Then

$$q_t(x) = x_2v_t + \tilde{q}_t(x)$$

where $\tilde{q}_t \in A_2(x_2, x_3)$ and $v_t \in A_1(x_1, x_2, x_3)$. Then equation 3.31 is written as

$$(A.1) \quad e^{\alpha t}((v_{t+s} - e^{\delta s}v_t)x_2 + (\tilde{q}_{t+s} - e^{\delta s}\tilde{q}_t)) = e^{\alpha t}x_2v_s(\ell_{it}) + \tilde{q}_s(\ell_{it}).$$

We denote $\tilde{q}_t = A(t)x_1^2 + 2B(t)x_1x_3 + C(t)x_3^2$ and $v_t(x) = a(t)x_1 + b(t)x_2 + c(t)x_3$. Then equation A.1 is written as

$$\begin{aligned} A(t+s) - e^{\delta s}A(t) &= e^{-\alpha t}A(s) \\ B(t+s) - e^{\delta s}B(t) &= e^{-\alpha t}(B(s) + tA(s)) \\ C(t+s) - e^{\delta s}C(t) &= e^{-\alpha t}(C(s) + 2tB(s) + t^2A(s)) \\ a(t+s) - e^{\delta s}a(t) &= a(s) + 2(1 - e^{-\alpha t})A(s) \\ b(t+s) - e^{\delta s}b(t) &= e^{\alpha t}b(s) + (e^{\alpha t} - 1)a(s) + (e^{\alpha t} - 1)(1 - e^{-\alpha t})A(s) \\ c(t+s) - e^{\delta s}c(t) &= c(s) + ta(s) + 2t(1 - e^{-\alpha t})A(s) + 2(1 - e^{-\alpha t})B(s), \end{aligned}$$

yielding (solving the equations corresponding to $a(t) + 2A(t)$, $a(t) + b(t) + A(t)$ and $c(t) + 2B(t)$):

$$\begin{aligned} A(t) &= A(e^{-\alpha t} - e^{\delta t}) \quad (\text{or } Ate^{\delta t} \quad \delta = -\alpha) \\ B(t) &= B(e^{-\alpha t} - e^{\delta t}) + Ate^{-\alpha t} \quad (\text{or } (Bt + \frac{A}{2}t^2)e^{\delta t} \quad \delta = -\alpha) \\ C(t) &= C(e^{-\alpha t} - e^{\delta t}) + 2Bte^{-\alpha t} + At^2e^{-\alpha t} \quad (\text{or } (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t} \quad \delta = -\alpha) \\ a(t) &= a(1 - e^{\delta t}) - 2A(e^{-\alpha t} - e^{\delta t}) \\ &\quad (\text{or } at - 2A(e^{-\alpha t} - 1) \quad \text{for } \delta = 0, \\ &\quad (\text{or } a(1 - e^{\delta t}) - 2Ate^{\delta t} \quad \text{for } \delta = -\alpha) \\ b(t) &= b(e^{\alpha t} - e^{\delta t}) - a(1 - e^{\delta t}) + A(e^{-\alpha t} - e^{\delta t}) \\ &\quad (\text{or } b(e^{\alpha t} - 1) - at + A(e^{-\alpha t} - 1) \quad \text{for } \delta = 0, \\ &\quad \text{or } b(e^{-\delta t} - e^{\delta t}) - a(1 - e^{\delta t}) + Ate^{\delta t} \quad \text{for } \delta = -\alpha, \\ &\quad \text{or } bte^{\delta t} - a(1 - e^{\delta t}) + Ate^{\delta t} \quad \text{for } \delta = \alpha) \\ c(t) &= c(1 - e^{\delta t}) + at - 2Ate^{-\alpha t} + 2B(1 - e^{-\alpha t}) \\ &\quad (\text{or } ct + \frac{a}{2}t^2 - 2Ate^{-\alpha t} + 2B(1 - e^{-\alpha t}) \quad \text{for } \delta = 0, \\ &\quad \text{or } c(1 - e^{\delta t}) + at - (2Bt + At^2)e^{\delta t} \quad \text{for } \delta = -\alpha), \end{aligned}$$

which yields flows

- $\varphi_t(x) = [e^{\delta t}x_0x_2 + (a(1 - e^{\delta t}) - 2A(e^{-\alpha t} - e^{\delta t}))x_1x_2 + (b(e^{\alpha t} - e^{\delta t}) - a(1 - e^{\delta t}) + A(e^{-\alpha t} - e^{\delta t}))x_2^2 + (c(1 - e^{\delta t}) + at - 2Ate^{-\alpha t} + 2B(1 - e^{-\alpha t}))x_2x_3 + A(e^{-\alpha t} - e^{\delta t})x_1^2 + 2(B(e^{-\alpha t} - e^{\delta t}) + Ate^{-\alpha t})x_1x_3 + (C(e^{-\alpha t} - e^{\delta t}) + At^2e^{-\alpha t} + 2Bte^{-\alpha t})x_3^2, (x_1 + (e^{\alpha t} - 1)x_2 + tx_3)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\delta \neq 0$, $\alpha \neq 0$,
- $\varphi_t(x) = [x_0x_2 + (at - 2A(e^{-\alpha t} - 1))x_1x_2 + (b(e^{\alpha t} - 1) - at + A(e^{-\alpha t} - 1))x_2^2 + (ct + \frac{a}{2}t^2 - 2Ate^{-\alpha t} + 2(1 - e^{-\alpha t})B)x_2x_3 + A(e^{-\alpha t} - e^{\delta t})x_1^2 + 2(B(e^{-\alpha t} - 1) + Ate^{-\alpha t})x_1x_3 + (C(e^{-\alpha t} - 1) + At^2e^{-\alpha t} + 2Bte^{-\alpha t})x_3^2, (x_1 + (e^{\alpha t} - 1)x_2 + tx_3)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\alpha \neq 0$,

- $\varphi_t(x) = [e^{\delta t}x_0x_2 - 2Ate^{\delta t}x_1x_2 + (b(e^{-\delta t} - e^{\delta t}) - a(1 - e^{\delta t}) + Ate^{\delta t})x_2^2 + (c(1 - e^{\delta t}) + at - 2Ate^{-\delta t} + (2Bt + At^2)e^{\delta t})x_2x_3 + Ate^{\delta t}x_1^2 + (2Bt + At^2)e^{\delta t}x_1x_3 + (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t}x_3^2, (x_1 + (e^{\delta t} - 1)x_2 + tx_3)x_2, e^{\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_2 + (a(1 - e^{\delta t}) - 2A(e^{-\delta t} - e^{\delta t}))x_1x_2 + (bte^{\delta t} - a(1 - e^{\delta t}) + A(e^{-\delta t} - e^{\delta t}))x_2^2 + (c(1 - e^{\delta t}) + at - 2Ate^{-\delta t} + 2B(1 - e^{-\delta t}))x_2x_3 + A(e^{-\delta t} - e^{\delta t})x_1^2 + 2(B(e^{-\delta t} - e^{\delta t}) + Ate^{-\delta t})x_1x_3 + (C(e^{-\delta t} - e^{\delta t}) + At^2e^{-\delta t} + 2Bte^{-\delta t})x_3^2, (x_1 + (e^{\delta t} - 1)x_2 + tx_3)x_2, e^{\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$.

iv) One can assume that $\alpha \neq 0$ for otherwise we are in case i) for $\ell = x_3$. We obtain the equations:

$$\begin{aligned}
A(t+s) - e^{\delta s}A(t) &= e^{\alpha t}A(s) \\
B(t+s) - e^{\delta s}B(t) &= B(s) \\
C(t+s) - e^{\delta s}C(t) &= e^{-\alpha t}C(s) \\
a(t+s) - e^{\delta s}a(t) &= e^{\alpha t}a(s) + 2te^{\alpha t}A(s) \\
b(t+s) - e^{\delta s}b(t) &= e^{\alpha t}b(s) + te^{\alpha t}a(s) + t^2e^{\alpha t}A(s) \\
c(t+s) - e^{\delta s}c(t) &= c(s) + 2tB(s).
\end{aligned}$$

Therefore:

$$\begin{aligned}
A(t) &= A(e^{\alpha t} - e^{\delta t}) \quad (\text{or } Ate^{\delta t} \text{ for } \delta = \alpha) \\
B(t) &= B(1 - e^{\delta t}) \quad (\text{or } Bt \text{ for } \delta = 0) \\
C(t) &= C(e^{-\alpha t} - e^{\delta t}) \quad (\text{or } Cte^{\delta t} \text{ for } \delta = -\alpha) \\
a(t) &= a(e^{\alpha t} - e^{\delta t}) + 2Ate^{\alpha t} \quad (\text{or } (at + At^2)e^{\delta t} \text{ for } \delta = \alpha) \\
b(t) &= b(e^{\alpha t} - e^{\delta t}) + ate^{\alpha t} + At^2e^{\alpha t} \quad (\text{or } (bt + \frac{a}{2}t^2 + \frac{A}{3}t^3)e^{\delta t} \text{ for } \delta = \alpha) \\
c(t) &= c(1 - e^{\delta t}) + 2tB \quad (\text{or } ct + Bt^2 \text{ for } \delta = 0),
\end{aligned}$$

which yield the flows

- $\varphi_t(x) = [e^{\delta t}x_0x_2 + (a(e^{\alpha t} - e^{\delta t}) + 2Ate^{\alpha t})x_1x_2 + (b(e^{\alpha t} - e^{\delta t}) + ate^{\alpha t} + At^2e^{\alpha t})x_2^2 + (c(1 - e^{\delta t}) + 2tB)x_2x_3 + A(e^{\alpha t} - e^{\delta t})x_1^2 + 2B(1 - e^{\delta t})x_1x_3 + C(e^{-\alpha t} - e^{\delta t})x_3^2, e^{\alpha t}(x_1 + tx_2)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- $\varphi_t(x) = [x_0x_2 + (a(e^{\alpha t} - 1) + 2Ate^{\alpha t})x_1x_2 + (b(e^{\alpha t} - 1) + ate^{\alpha t} + At^2e^{\alpha t})x_2^2 + (ct + Bt^2)x_2x_3 + A(e^{\alpha t} - 1)x_1^2 + 2Btx_1x_3 + C(e^{-\alpha t} - 1)x_3^2, e^{\alpha t}(x_1 + tx_2)x_2, e^{\alpha t}x_2^2, x_2x_3]$ with $\alpha \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_2 + (at + At^2)e^{\delta t}x_1x_2 + (bt + \frac{a}{2}t^2 + \frac{A}{3}t^3)e^{\delta t}x_2^2 + (c(1 - e^{\delta t}) + 2tB)x_2x_3 + Ate^{\delta t}x_1^2 + 2B(1 - e^{\delta t})x_1x_3 + C(e^{-\delta t} - e^{\delta t})x_3^2, e^{\delta t}(x_1 + tx_2)x_2, e^{\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_2 + (a(e^{-\delta t} - e^{\delta t}) + 2Ate^{-\delta t})x_1x_2 + (b(e^{-\delta t} - e^{\delta t}) + ate^{-\delta t} + At^2e^{-\delta t})x_2^2 + (c(1 - e^{\delta t}) + 2tB)x_2x_3 + A(e^{-\delta t} - e^{\delta t})x_1^2 + 2B(1 - e^{\delta t})x_1x_3 + Cte^{\delta t}x_3^2, e^{-\delta t}(x_1 + tx_2)x_2, e^{-\delta t}x_2^2, x_2x_3]$ with $\delta \neq 0$.

3) $\ell = x_1$ and Ψ_t of type:

i) Then $\lambda(t) = e^{\alpha t}$ and

$$q_t(x) = x_1v_t + \tilde{q}_t(x)$$

where $\tilde{q}_t \in A_2(x_2, x_3)$ and $v_t \in A_1(x_1, x_2, x_3)$. Equation 3.31 is written as

$$(A.2) \quad x_1(v_{t+s} - e^{\delta s}v_t) + (\tilde{q}_{t+s} - e^{\delta s}\tilde{q}_t) = x_1v_s(\ell_{it}) + e^{-\alpha t}\tilde{q}_s(\ell_{it}).$$

We denote $\tilde{q}_t = A(t)x_2^2 + 2B(t)x_2x_3 + C(t)x_3^2$ and $v_t(x) = a(t)x_1 + b(t)x_2 + c(t)x_3$. Then equation A.1 is written as

$$\begin{aligned}
A(t+s) - e^{\delta s}A(t) &= e^{-\alpha t}A(s) \\
B(t+s) - e^{\delta s}B(t) &= e^{-\alpha t}(B(s) + tA(s)) \\
C(t+s) - e^{\delta s}C(t) &= e^{-\alpha t}(C(s) + 2tB(s) + t^2A(s)) \\
a(t+s) - e^{\delta s}a(t) &= e^{\alpha t}a(s) \\
b(t+s) - e^{\delta s}b(t) &= b(s) \\
c(t+s) - e^{\delta s}c(t) &= c(s) + tb(s).
\end{aligned}$$

with solutions:

$$\begin{aligned}
 A(t) &= A(e^{-\alpha t} - e^{\delta t}) \quad (\text{or } Ate^{\delta t} \text{ for } \delta = -\alpha) \\
 B(t) &= B(e^{-\alpha t} - e^{\delta t}) + Ate^{-\alpha t} \quad (\text{or } (Bt + \frac{A}{2}t^2)e^{\delta t} \text{ for } \delta = -\alpha) \\
 C(t) &= C(e^{-\alpha t} - e^{\delta t}) + 2Bte^{-\alpha t} + At^2e^{-\alpha t} \quad (\text{or } (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t} \text{ for } \delta = -\alpha) \\
 a(t) &= a(e^{\alpha t} - e^{\delta t}) \quad (\text{or } ate^{\delta t} \text{ for } \alpha = \delta) \\
 b(t) &= b(1 - e^{\delta t}) \quad (\text{or } bt \text{ for } \delta = 0) \\
 c(t) &= c(1 - e^{\delta t}) + bt \quad (\text{or } ct + \frac{b}{2}t^2 \text{ for } \delta = 0),
 \end{aligned}$$

which yield the flows

- $\varphi_t(x) = [e^{\delta t}x_0x_1 + a(e^{\alpha t} - e^{\delta t})x_1^2 + b(1 - e^{\delta t})x_1x_2 + (c(1 - e^{\delta t}) + tb)x_1x_3 + A(e^{-\alpha t} - e^{\delta t})x_2^2 + 2(B(e^{-\alpha t} - e^{\delta t}) + Ate^{-\alpha t})x_2x_3 + (C(e^{-\alpha t} - e^{\delta t}) + (At^2 + 2tB)e^{-\alpha t})x_3^2, e^{\alpha t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_1 + a(e^{-\delta t} - e^{\delta t})x_1^2 + b(1 - e^{\delta t})x_1x_2 + (c(1 - e^{\delta t}) + tb)x_1x_3 + Ate^{\delta t}x_2^2 + 2(Bt + \frac{A}{2}t^2)e^{\delta t}x_2x_3 + (Ct + Bt^2 + \frac{A}{3}t^3)e^{\delta t}x_3^2, e^{-\delta t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\delta \neq 0$,
- $\varphi_t(x) = [e^{\delta t}x_0x_1 + ate^{\delta t}x_1^2 + b(1 - e^{\delta t})x_1x_2 + (c(1 - e^{\delta t}) + tb)x_1x_3 + A(e^{-\delta t} - e^{\delta t})x_2^2 + 2(B(e^{-\delta t} - e^{\delta t}) + Ate^{\delta t})x_2x_3 + (C(e^{-\delta t} - e^{\delta t}) + (At^2 + 2tB)e^{-\delta t})x_3^2, e^{\delta t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\delta \neq 0$,
- $\varphi_t(x) = [x_0x_1 + a(e^{\alpha t} - 1)x_1^2 + btx_1x_2 + (ct + \frac{b}{2}t^2)x_1x_3 + A(e^{-\alpha t} - 1)x_2^2 + 2(B(e^{-\alpha t} - 1) + Ate^{-\alpha t})x_2x_3 + (C(e^{-\alpha t} - 1) + (At^2 + 2tB)e^{-\alpha t})x_3^2, e^{\alpha t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$ with $\alpha \neq 0$,
- $\varphi_t(x) = [x_0x_1 + atx_1^2 + btx_1x_2 + (ct + \frac{bt^2}{2})x_1x_3 + Atx_2^2 + (2Bt + At^2)x_2x_3 + (Ct + Bt^2 + \frac{A}{3}t^3)x_3^2, e^{\alpha t}x_1^2, (x_2 + tx_3)x_1, x_3x_1]$.

REFERENCES

1. M. Alberich-Carramiñana, *Geometry of the plane Cremona maps*, Lecture Notes in Mathematics, vol. 1769, Springer-Verlag, Berlin, 2002.
2. L. Bayle and A. Beauville, *Birational involutions of $\mathbf{P}^2$* , Asian J. Math. **4** (2000), no. 1, 11–17, Kodaira's issue.
3. E. Bertini, *Richerche sulle trasformazioni univoche involutorie nel piano*, Annali di Mat. **8** (1877), 244–286.
4. J. Blanc, I. Pan, and T. Vust, *Sur un théorème de Castelnuovo*, Bull. Braz. Math. Soc. **39**, no. 1.
5. S. Cantat and C. Favre, *Symétries birationnelles des surfaces feuilletées*, J. Reine Angew. Math. **561** (2003), 199–235.
6. G. Castelnuovo, *Sulle trasformazioni cremoniane del piano, che ammettono una curva fiera*, Rend. Accad. Lincei (1892).
7. D. Cerveau and J. Déserti, *Transformations birationnelles de petit degré*, arxiv:0811.2325[math.AG].
8. M. Demazure, *Sous-groupes algébriques de rang maximum du groupe de Cremona*, Ann. Sci. École Norm. Sup. (4) **3** (1970), 507–588.
9. J. Déserti, *Groupe de cremona et dynamique complexe: une approche de la conjecture de Zimmer*, Int. Math. Res. Not. **Art. ID 71701**. (2006).
10. I.V. Dolgachev and V. A. Iskovskih, *Groupe de cremona et dynamique complexe: une approche de la conjecture de Zimmer*, Int. Math. Res. Not. **Art. ID 71701**. (2006).
11. F. Enriques and G. Fano, *Sui gruppi di trasformazioni cremoniane dell spazio*, Annali di Matematica pura ed applicata, s. 2a **15** (1897), 221–271.
12. G. Fano, *I gruppi di Jonquières generalizzati*, Atti della R. Acc. di Torino **33** (1898), 59–98.
13. L. Godeaux, *Les transformations birationnelles du plan*, Mémorial des Sciences Mathématiques, vol. 122, Gauthier-Villars, Paris, 1953.
14. H. Hudson, *Cremona transformations in plane and space*, Cambridge University Press, London, 1927.
15. V. A. Iskovskih, *Proof of a theorem on relations in the two-dimensional cremona group*, Uspekhi Mat. Nauk. **40** (1985), no. 5(245), 255–256.
16. S. Kantor, *Theorie der endlichen gruppen von eindeutigen transformationen in der ebene*, Mayer & Mullen, 1895.
17. I. Pan, *Une remarque sur la génération du groupe de Cremona*, Bol. Soc. Brasil. Mat. (N.S.) **30** (1999), no. 1, 95–98.
18. I. Pan, F. Ronga, and T. Vust, *Transformations birationnelles quadratiques de l'espace projectif complexe à trois dimensions*, Ann. Inst. Fourier (Grenoble) **51** (2001), no. 5, 1153–1187.
19. H. Umemura, *Maximal algebraic subgroups of the Cremona group of three variables. Imprimitve algebraic subgroups of exceptional type*, Nagoya Math. J. **87** (1982), 59–78.
20. ———, *On the maximal connected algebraic subgroups of the Cremona group. I*, Nagoya Math. J. **88** (1982), 213–246.
21. ———, *On the maximal connected algebraic subgroups of the Cremona group. II*, Algebraic groups and related topics (Kyoto/Nagoya, 1983), Adv. Stud. Pure Math., vol. 6, North-Holland, Amsterdam, 1985, pp. 349–436.